COUNTWAY LIBRARY
HC 52BV Z

BOSTON
MEDICAL LIBRARY
8 THE FENWAY

Medical and Surgical
JOURNAL.

RETINOSCOPY

(SHADOW TEST)

THORINGTON

Reviews of Previous Editions of

Thorington's Retinoscopy.

From "The Medical Record," New York.

"This little manual deserves a second edition, and will undoubtedly pass through many more. It presents a clear, terse, and thorough exposition of an objective method of determining refraction errors which is deservedly increasing in popularity. In our opinion the author is amply justified in declaring that its great value in nystagmus, young children, amblyopia, aphakia, and in examining illiterates and the feeble-minded can not be overestimated, and we agree with him in reminding those who attempt retinoscopy, fail, and ridicule it, that the fault is behind and not in front of the mirror. The book is well printed and usefully illustrated."

From "The Annals of Ophthalmology," St. Louis, Mo.

"Retinoscopy has come to stay. It is not a fad, neither a fashion. It is scientific-and withal so eminently practical in its application as to commend it to every think, ing worker in ophthalmology. The tendency in the medicine of to-day is toward objective methods. An objective method must possess two attributes: exactness and absolute independence of the patient's testimony. In addition to these qualities, an objective method must, if it is to meet with general acceptance, be easy of application. Ophthalmoscopy and ophthalmometry are but relatively exact in refractive work, seeing which the trial-case has held its supremacy up to date; nor would we wish to relegate it to the background. With a patient whose testimony is trustworthy exact results are thus obtainable, but it requires the most intelligent cooperation on the part of the examined. If, however, there be but the least departure from the conditions essential to close work with the test-lenses,—as, for instance, with foreigners, illiterates, children, partial amblyopics, or mental astigmatics,—retinoscopy stands ready to furnish a verdict from which there can be no appeal, when one has learned to properly interpret the movements observed in the pupillary area. It is to the elucidation of these latter movements as observed through a plane mirror at a distance of one meter that Dr. Thorington devotes himself in the volume before us. The treatment of the subject is so beautifully simple that one who runs may read."

From "The Journal of the American Medical Association," Chicago, Ill.

"The author of this well-written little book has very satisfactorily described the most approved methods of retinoscopy. The work is especially valuable in that for a great part it details the results of personal investigation of so well-known an authority on this subject as Dr. Thorington. Oculists accustomed to casually use retinoscopy as practised in the old way, with the concave mirror or with the ophthalmoscopic mirror, will be surprised to note the marked evolution of the modus operandi of this test as developed by Drs. Jackson and Thorington. With perfected instruments and strict attention to arrangement of light, distance, and other details, a surprising degree of proficiency and accuracy is possible. Any one pursuing the modern methods of retinoscopy will soon be convinced of its superiority over all other objective tests, and every worker in ophthalmology realizes the necessity of at least one reliable objective method of refraction."

From "The New Orleans Medical and Surgical Journal," New Orleans, La.

"We have nothing but a good word for this little book. It seems to fulfil well the purpose intended. It gives a brief, clear description of the means and manner of retinoscopy, together with the principles or natural laws upon which it is founded. The author has done well in selecting the method he thinks best and simplest, and has confined himself to it, so that the student will have no difficulty or confusion in following the manual step by step, and learning to put in practice for himself what is described in the pages. This once accomplished, he can readily, if he becomes convinced of its usefulness, acquire the variations and refinement upon this mode of examination."

From "The New York Medical Journal," New York.

"This little book presents as simple and practical a description of the shadow test as exists in our language."

From "The Scottish Medical and Surgical Journal," Edinburgh, Scotland.

"Dr. Thorington's lucid text is accompanied by twenty-four good illustrations, and on every page one notes that careful attention has been paid to little details of manipulation which stamp the writer as a practical teacher."

From "The Homeopathic Eye, Ear, and Throat Journal," New York.

"A practical and useful book. This is one of the most concise and clearest explanations of this subject we have seen. Retinoscopy is one of the most valuable aids we have in refractive work."

From "The Denver Medical Times," Denver, Col.

"His directions and descriptions are exceptionally clear and concise, and the little book he has written, we think, will be helpful to every physician who is interested in the fitting of glasses."

From "The Chicago Medical Recorder," Chicago, Ill.

"This little book is the most practical and complete exposition of the value and application of the shadow test in determining refractive errors with which we have any acquaintance. The illustrations, directions, advice, and general information in the book are all admirable."

From "The Post-Graduate," New York.

"This work on retinoscopy is divided into six chapters and an index. As stated in the preface, it is an abstract of the author's previous writings and lectures on retinoscopy, delivered at the Philadelphia Polyclinic. It is intended for college students and post-graduates, yet it is sufficiently complete for the use of the ophthalmologist. Retinoscopy has been selected as the name of the test, as it is the retina in its relative position to the refractive media which is studied. Skiascopy and skiagraphy are therefore regarded as misleading. To all those who are interested in this test for the determination of refraction we commend the work."

From "The Philadelphia Polyclinic," Philadelphia.

"We take pleasure in commending this concise statement of the methods to be employed in the routine use of a most valuable objective means of determining the errors of refraction. The student is told in simple English how to proceed in the examination."

From "The Boston Medical and Surgical Journal," Boston, Mass.

"This little manual is certainly the clearest exposition of this method of estimating refraction of the eye that has yet been published. The methods described are not so complicated as those taught in some other handbooks. The text is clear, and the illustrations serve the purposes for which they are designed admirably. Taken altogether, it is the most practicable handbook on retinoscopy yet published."

From "The Journal of Ophthalmology, Otology, and Laryngology," New York.

"We most emphatically recommend this little book to the beginner in the study of this method of determining refraction. The title is an index of the character of the text. It is positive, exact, practical. The aim of the author has been to present facts, and in as small space as possible. He has succeeded absolutely. The average work on this subject is, to the beginner, somewhat confusing, from the amount of theory presented—theory which is not always clear to the student. This has been avoided in the present case. Little, if any, theory is included, and the monograph is a series of categorical statements—clear, precise, and sufficient."

*** The price of this book (fourth edition, revised and enlarged, with eight additional illustrations) is $1.00 net, upon receipt of which it will be sent postpaid to any address. It may be obtained from the publishers or through any bookseller or dealer in opticians' supplies.

BY THE SAME AUTHOR

REFRACTION AND HOW TO REFRACT. With 200 Illustrations, many of which are from Original Drawings, Thirteen being Colored. Second Edition. *Just Ready.* Cloth, net $1.50

"It is a sterling book."—*Annals of Ophthalmology.*

"There is not a word of trash from cover to cover; everything is concise, accurate, sufficient, and up to date."—*The Journal of Ophthalmology, Otology and Laryngology.*

"The book can be recommended not only to beginners in the study of ophthalmology, but to those practitioners and students as well, whose limited knowledge of mathematics precludes the study of Helmholtz or Donders."—*New York Medical Record.*

"The many formulæ which are frequently brought into the text serve very well to elucidate the problems therein presented."—*Boston Medical and Surgical Journal.*

"It is clearly written and very fully illustrated, and will furnish an aid to the understanding of much that is often practically Greek to others than specialists."—*Journal of the American Medical Association, Chicago.*

"It will be found extremely useful to students, and will also be of great use to practitioners who want a handy book of reference, and one which is not overcrowded with minute details which are of theoretical interest only."—*British Medical Journal, London.*

P. Blakiston's Son & Co., Publishers, Philadelphia

RETINOSCOPY

(OR SHADOW TEST)

IN THE

DETERMINATION OF REFRACTION AT ONE METER DISTANCE, WITH THE PLANE MIRROR

BY

JAMES THORINGTON, A.M., M.D.

AUTHOR OF "REFRACTION AND HOW TO REFRACT"; PROFESSOR OF DISEASES OF THE EYE IN THE PHILADELPHIA POLYCLINIC AND COLLEGE FOR GRADUATES IN MEDICINE; OPHTHALMOLOGIST TO THE ELWYN AND VINELAND TRAINING SCHOOLS FOR FEEBLE-MINDED CHILDREN; OPHTHALMOLOGIST TO THE M. E. ORPHANAGE, ETC.

FOURTH EDITION, REVISED AND ENLARGED

FIFTY-ONE ILLUSTRATIONS

TWELVE OF WHICH ARE COLORED

PHILADELPHIA

P. BLAKISTON'S SON & CO.

1012 WALNUT STREET

1901

20130

COPYRIGHT, 1901, BY JAMES THORINGTON, M.D.

28. E. 39.



PRESS OF WM. F. FELL & CO.,
1220-24 SANSOM STREET,
PHILADELPHIA.

THIS BOOK IS AFFECTIONATELY DEDICATED TO THE MEMORY OF

FELIX A. BETTELHEIM, PH.D., M.D.,

MY FRIEND AND ASSOCIATE DURING HIS SIX YEARS' RESIDENCE, AS SURGEON OF THE PANAMA RAILROAD COMPANY, AT PANAMA.

PREFACE TO FOURTH EDITION.

The increasing demand for this small book is certainly very pleasing to the writer, especially as translations into the French and German have been asked for and are in course of preparation. In preparing this fourth edition of Retinoscopy the same endeavor has been made as formerly to present the facts to the reader in the same aspect which they bear to the author, as viewed in the light of his personal experience. To further this study, the whole subject has been carefully gone over, new paragraphs added, and eight new illustrations have been inserted.

120 S. Eighteenth St., Philadelphia, Pa.

January, 1901.

PREFACE TO THIRD EDITION.

The first edition of "Retinoscopy" was published in March, 1897, and it is most gratifying to the author that the subject continues to merit sufficient attention to call for a third edition in less than two years. Endeavoring to make the book more worthy of the favor with which it has been received and for a clearer understanding of the subject, some few changes in phraseology have been made and five additional illustrations incorporated.

120 S. Eighteenth St., Philadelphia, Pa.,
February, 1899.

PREFACE TO THE SECOND EDITION.

The first edition of this book was published in March, 1897, and it is indeed gratifying to the author that the work has found such favor as to call for a second in so short a time.

To make this edition more lucid than the first, the writer has carefully reviewed the original text and made some changes in the phraseology, and at the same time has added many new illustrations, twelve of which are in colors.

A description and drawings of three lenses, suggested by the author for the study of the scissor movement, conic cornea, and spheric aberration on the schematic eye, have also been inserted.

120 S. Eighteenth St., Philadelphia, Pa.,
February, 1898.

PREFACE TO THE FIRST EDITION.

At the earnest solicitation of many students and friends, this book is presented as an abstract of the author's previous writings and lectures on Retinoscopy, delivered during the winter course on Ophthalmology, at the Philadelphia Polyclinic.

In presenting a manual of this kind the writer does not presume to detract from the writings or teachings of others, or the excellent work on Skiascopy, by his friend and colleague, Dr. E. Jackson; but wishes to elucidate in as concise a manner and few words as possible the method of applying retinoscopy, which has given most satisfaction at his hands.

While intended for college students and post-graduates, yet there is ample material given whereby the ophthalmologist at a distance may acquire a working knowledge of the method, by study and practice in his own office.

For three reasons Retinoscopy, in preference to Skiascopy, has been chosen as the title:

First, that it may not be confounded with Skiagraphy.

Second, that it is the name by which the test is universally known; and—

Third, that it is the retina in its relative position to the dioptric media which we study.

120 S. Eighteenth St., Philadelphia, Pa.,
March, 1897.

CONTENTS.

CHAPTER I.

PAGE

Definition.—Names.—Principle and Value of Retinoscopy.—Suggestions to the Beginner, ... 19–23

CHAPTER II.

Retinoscope.—Light.—Light-screen.—Dark Room.—Source of Light and Position of Mirror.—Observer and Patient, . 24–30

CHAPTER III.

Distance of Surgeon from Patient.—Arrangement of Patient, Light, and Observer.—Reflection from Mirror.—How to Use the Mirror.—What the Observer Sees.—Retinal Illumination.—Shadow.—Where to Look and What to Look For, ... 31–39

CHAPTER IV.

Point of Reversal.—To Find the Point of Reversal.—What to Avoid.—Direction of Movement of Retinal Illumination.—Rate of Movement and Form of Illumination.—Rules for Lenses.—Movement of Mirror and Apparatus, 40–51

CHAPTER V.

Retinoscopy in Emmetropia and the Various Forms of Regular Ametropia.—Axonometer, ... 52–69

CHAPTER VI.

Retinoscopy in the Various Forms of Irregular Ametropia.—Retinoscopy without a Cycloplegic.—The Concave Mirror.—Description of the Author's Schematic Eye and Light-screen. — Lenses for the Study of the Scissor Movement, Conic Cornea, and Spheric Aberration, . . . 70–86

INDEX, ... 87–89

LIST OF ILLUSTRATIONS.

FIG.		PAGE
1.	Schematic Eye for Studying Retinoscopy,	22
2.	Retinoscope,	25
3.	Oil Lamp,	26
4.	Light-screen, or Cover Chimney,	26
5.	New Light-screen,	27
6.	Showing Distance from Patient's Eyes and the Equivalent in Diopters,	32
7.	Arrangement of Patient, Light, and Observer,	33
8.	Light over Patient's Head, and the Observer with Mirror at One Meter Distance,	34
9.	Folding Mirror,	35
10.	Folding Mirror with Illumination,	35
11.	Illumination in an Emmetropic Eye,	38
12.	Illumination and Shadow in an Emmetropic Eye,	38
13.	Illumination with Straight Edge,	46
14.	Illumination with Crescent Edge,	46
15.	Würdemann's Disc,	47
16.	Jennings' Skiascopic Disc,	48
17.	Trial Frame,	50
18.	Trial-case,	51
19.	Gray Reflex as seen in High Hyperopia,	52
20.	Gray Reflex, Crescent Edge, and Shadow in High Hyperopia,	52
21.	Hyperopia,	53
22.	Refracted Hyperopia,	54
23.	Emmetropia,	55
24.	Refraction of Macular Region,	56
25.	Myopia,	57
26.	Refracted Myopia,	58
27.	Method of Writing a Formula,	61
28.	Band of Light in Astigmatism,	62
29.	Band of Light and Shadow,	62
30.	Band of Light, Axis 90°,	63
31.	Band of Light Showing Half a Diopter of Astigmatism,	64
32.	Axonometer,	67
33.	Axonometer in Position,	67

FIG. PAGE
34. Celluloid Axonometer, ... 69
35, 36. Irregular Lenticular Astigmatism, ... 71
37. Two Bands of Light, ... 73
38. Light Areas with Dark Interspace, ... 72
39. Light Areas Brought Together, ... 73
40. Tilting of Lens, ... 74
41. Scissor Movement in Refracted Aphakia, ... 75
42. Illumination Seen in Conic Cornea, ... 77
43. Positive Aberration, ... 78
44. Negative Aberration, ... 78
45. Lens for the Study of the Scissor Movement, ... 83
46. Lens for Study of the Scissor Movement, ... 84
47. Lens for the Study of Conic Cornea, ... 83
48. Lens for Study of Conic Cornea, ... 84
49. Lens for the Study of Spheric Aberration, ... 83
50. Lens for Study of Spheric Aberration, ... 85
51. Lens for Study of Irregular Lenticular Astigmatism, ... 85

FEB 15

RETINOSCOPY.

CHAPTER I.

DEFINITION.—NAMES.—PRINCIPLE AND VALUE OF RETINOSCOPY.—SUGGESTIONS TO THE BEGINNER.

Definition.—Retinoscopy (see preface to the first edition) may be defined as the method of estimating the refraction of an eye by reflecting into it rays of light from a plane or concave mirror, and observing the movement which the retinal illumination makes by rotating the mirror.

Names.—Dioptroscopy, fundus-reflex test, keratoscopy, fantoscopy, pupilloscopy, retinophotoscopy, retinoskiascopy, skiascopy, umbrascopy, koroscopy, etc., are some of the other names given to this method of estimating the refraction, and their number and greater or less inappropriateness have had much to do, no doubt, with keeping retinoscopy in the background of ophthalmology instead of giving it the prominence which it more justly deserved and is now receiving from ophthalmologists in all parts of the world.

The principle of retinoscopy is the finding of the point of reversal (the far-point of a myopic eye), and to do this, if an eye is not already sufficiently myopic, it will be necessary to place in front of it such a lens, or series of lenses, as will bring the emergent rays of light to a focus at a certain definite distance (see Point of Reversal, chap. IV).

Value of Retinoscopy.—Those who would criticize retinoscopy because "we see nothing and think nothing of the condition of the fundus," base their criticism apparently on the name, retinoscopy, rather than from any great amount of practical experience with the method. While admitting that the ophthalmoscope in front of a well-trained eye will often give a close refractive estimate of the eye under examination, yet only to the few does such skill obtain, and even then there is that uncertainty which does not attach itself to the retinoscope in competent hands. The ophthalmologist who knows how to use the retinoscopic mirror accurately has the advantage of his confrères who are ignorant of the test; it gives him a position decidedly independent of his patient, and puts him above the common level of the traveling "Great Doctor Eye" and "refracting optician," who are tied to the trial-lenses and the patient's uncertain answers. Furthermore, when it is remembered that from fifty to eighty per cent. of the patients consulting the ophthalmologist do so for an error of refraction, it is well that he be most capable in this important branch of the subject.

The wonderful advantage of retinoscopy over other methods needs no argument to uphold it; the

rapidly increasing number of retinoscopists testify to its merits.

The writer, from his constant use of the mirror, would suggest the following axiom: That, *with an eye otherwise normal except for its refractive error, and being under the influence of a reliable cycloplegic, there is no more accurate objective method of obtaining its exact correction than by retinoscopy.*

Retinoscopy gives the following advantages:

The character of the refraction is quickly diagnosed.

The exact refraction is obtained without questioning the patient.

Little time is required to make the test.

No expensive apparatus is necessarily required.

Its great value can never be overestimated in cases of nystagmus, young children, amblyopia, aphakia, illiterates, and the feeble-minded.

From what has just been written, it must not be understood that the patient's glasses are ordered immediately from the result obtained by retinoscopy; for, on the contrary, all retinoscopic work, like ophthalmometry in general, should, when possible, be confirmed at the trial-case.

It is only in the feeble-minded, in young children, and in cases of amblyopia that glasses are ordered direct from the findings obtained in the dark room.

The subjective method of placing lenses before the patient's eyes and letting him decide by asking "is this better?" or "is this worse?" only too often fatigues the examiner and worries the patient, giving him or her a dread or fear of inaccuracy that does

not satisfy the surgeon or tend to inspire the patient. Whereas, when the neutralizing lenses found by retinoscopy are placed before the patient's eyes and he reads $\frac{6}{6}$ or $\frac{20}{20}$ or more, it is easy, if there is any doubt, to hold up a plus and a minus quarter diopter

FIG. 1.—THE AUTHOR'S SCHEMATIC EYE FOR STUDYING RETINOSCOPY. (*For description, see chap. vi.*)

glass respectively in front of this correction, and let the patient tell at once if either glass improves or diminishes the vision.

The writer is not condemning the subjective or other methods of estimating the refraction, nor is

he trying to extol too highly the shadow test, yet he would remind those who try retinoscopy, fail, and then ridicule it, that the fault with them is *back* and not in front of the mirror.

Suggestions to the Beginner.—To obtain proficiency in retinoscopy there is much to be understood. Careful attention to details *must* be given, and not a little patience possessed, as it is not a method that is acquired in a day, and it is only after weeks of constant application that accuracy is acquired. Therefore the beginner is strongly advised to learn the major points from one of the many schematic eyes in the market before attempting the human eye. At the same time he should be perfectly familiar with the laws of refraction and dioptrics, as an understanding of conjugate foci is really the underlying principle of the method—*i. e.*, a point on the retina being one focus and the myopic or artifically-made far-point the other focus.

What is meant by major points applies more particularly to the study of the retinal illumination, its direction and apparent rate of movement, also its form, the distance between the observer and the patient, how to handle the mirror, etc., all of which are referred to under their special headings.

CHAPTER II.

RETINOSCOPE.—LIGHT.—LIGHT-SCREEN.—DARK ROOM.—SOURCE OF LIGHT AND POSITION OF MIRROR.—OBSERVER AND PATIENT.

The Retinoscope, or Mirror.—Two forms of the plane mirror are in use—the one large, four centimeters in diameter, with a four- or five-millimeter sight-hole often cut through the glass; and the other small, two centimeters in diameter, on a four-centimeter metal disc, with sight-hole two millimeters in diameter, *not* cut through the glass, the quicksilver or plating alone being removed. By thus leaving the glass at the sight-hole, additional reflecting surface is obtained at this point, which assists materially in exact work, as it diminishes the dark central shadow that shows so conspicuously at times, and particularly when the sight-hole is cut through the glass. The small mirror has an advantage over the large by reducing the area of reflected light, as only a one-centimeter area on each side of the sight-hole is of particular use. The small plane mirror* is the one recommended, and is made with either a straight or folding handle (see Fig. 10); the latter is for the purpose of protecting the mirror when carried in the pocket. The

**Philadelphia Polyclinic*, November, 1893. Another form is described by Dr. E. Jackson, *American Journal of Ophthalmology*, April, 1896.

purpose of the metal disc on which the small mirror is secured is to keep the light out of the observer's eye, and enable him to rest the instrument against the brow and side of the nose; but if its size should appear small, the observer can easily have a larger one made to suit his convenience. The plating or silvering on the mirror should be of the best, and free from any flaws or imperfections, for on its quality depends, in part, the good reflecting power of the mirror, which is very important.



FIG. 2.—THE AUTHOR'S RETINOSCOPE.*

The central shadow just referred to as the result of the sight-hole had best be seen by the beginner by reflecting the light from the mirror onto a white surface, before he begins any study, as this dark area may annoy him later if he does not understand its origin.

The Light.—This should be steady, clear, and white. The Welsbach possesses all these qualities, but unfortunately its delicate mantle will not stand

* See foot-note on preceding page.

much jarring, and as a consequence, is easily broken, causing much loss of time and annoyance. The electric light made with a twisted carbon and ground-glass covering having a round center of clear glass is becoming quite popular. For constant service, however, the Argand burner is decidedly the best, when the asbestos light-screen is used to intercept the heat. Whatever light is employed, it is well to have it on an extension bracket, so that the observer



FIG. 3.

may raise or lower it or move it toward or away from the patient, as necessary.

When gas or electric light is not at hand, a student's oil-lamp, with a suitable light-screen, will answer every purpose (see Fig. 3).

The light-screen, or cover chimney, is made of one-eighth inch asbestos, and of sufficient size (six centimeters in diameter by twenty-one in height) to fit over the glass chimney of the Argand burner (see Fig. 4).

Attached to the screen are two superimposed revolving discs that furnish four round openings, respectively five, ten, twenty, and thirty millimeters, any one of which may be turned into place as occasion may require. Care should be taken that the opening used is placed opposite to the brightest, and never opposite to the edge or the blue part of the



FIG. 4.—THE AUTHOR'S LIGHT-SCREEN, OR COVER CHIMNEY.
(*For a further description, see chap. vi.*)

FIG. 5.—THE AUTHOR'S NEW LIGHT-SCREEN.

flame. Formerly these screens were made of sheet-iron, but the asbestos has been found preferable, as it does not radiate the heat to the same extent as the iron. The purpose of the light-screen is to cover all of the flame except the portion which presents at the opening in the disc.

Ten-millimeter Opening.—This will be used in most all retinoscopic work by the beginner.

Five-millimeter Opening.—This is used to the best advantage and with no small amount of satisfaction by the expert when working close to the point of reversal. Figure 5 shows the author's new light-screen, which was described on page 1378 in the "Journal of the American Association," December 3, 1898. This is a more convenient screen for retinoscopy than the one shown in figure 4. It is made by attaching an iris diaphragm to an asbestos chimney. The amount of light passing through the diaphragm is easily controlled by an ivory-tipped lever at the left-hand side; and an index on the periphery records the diameter of the opening in use, from one to thirty millimeters.

The room must be darkened,—and the darker the better; all other sources of light except the one in use should be excluded. It must not be supposed from this that the room must have its walls and ceiling blackened; on the contrary, if the shades are drawn, the room will be sufficiently dark, though of course a perfectly black room would be best, as giving a greater contrast to the condition to be studied. The exclusion of other lights, or beams of light, must be insisted upon, as the principal use of the darkened room is to keep all light except the light in use out of the eye to be examined, and also not to have other lights reflected from the mirror.

As the method of using the concave mirror with source of light (twenty or thirty mm. opening in screen) beyond its principal focus (usually over and beyond the patient's head) has been superseded by the simpler and easier method of using the small plane

mirror with source of light (one-half or one cm. opening in light-screen) brought as close to the mirror as possible, the description of retinoscopy which follows will refer to the latter.

The Source of Light and Position of the Mirror.—The rays of light coming out of the round opening in the light-screen should be five or six inches to the left and front of the observer, so that they may pass *in front* of the left eye and fall upon the mirror held before the right, thus leaving the observer's left eye in comparative darkness; or the observer may use the mirror before the left eye in case he is left-handed and has the light to his right. It is always best for the observer to keep both eyes wide open and to avoid having any light fall into the unused eye, which would cause him much annoyance. Some observers hold the mirror before the eye next to the screen, but this is not recommended, for the reasons just mentioned.

The observer need not make any note of his accommodation, as in using the ophthalmoscope, but, as he requires very acute vision, he should wear any necessary correcting glasses. *Any observer whose vision does not approximate* $\frac{6}{6}$ *in the eye which he uses will not get much satisfaction from retinoscopy.*

He should take his seat facing the patient, and, as the strength or brilliancy of the reflected light rapidly weakens as the distance between the mirror and the light-screen is increased, he should have the light-screen close to his face (not farther away than six inches) if he wishes to get the fullest possible strength of light on the mirror.

As the light appears just as far back in the mirror as it is in front of it, then the nearer these two objects are brought together, the more nearly do they become as one. When working close to the point of reversal, more exact work will be accomplished if this distance between the light and mirror is very short. The nearer together the light and mirror, the brighter the retinal illumination, and greater contrast, or sharper cut edge between illumination and surrounding shadow. The further the light from the mirror, the dimmer the retinal illumination, and there will appear, under certain conditions, a very conspicuous central shadow as the result of the sight-hole in the mirror—*two very serious objections.*

The patient *must have his accommodation thoroughly relaxed with a reliable cycloplegic*, and should be seated comfortably, one meter distant, in front of the observer, with his vision steadily fixed on the observer's forehead, just above the mirror. Or, what is even better, the patient may concentrate his vision on the edge of the metal disc of the mirror or on the observer's forehead, but never directly into the mirror, as that would soon irritate and compel him to close his eye.

In this way the patient avoids the strain of looking into the bright reflected light, and at the same time the *macular region* is refracted (see Fig. 24). It is customary to cover the patient's *other* eye while its fellow is being refracted; for obvious reasons this is specially important in cases of "squint." The axonometer placed before the eye being refracted is a decided advantage in any instance (see p. 69).

CHAPTER III.

DISTANCE OF SURGEON FROM PATIENT.—ARRANGEMENT OF PATIENT, LIGHT, AND OBSERVER—REFLECTION FROM MIRROR.—HOW TO USE THE MIRROR.—WHAT THE OBSERVER SEES.—RETINAL ILLUMINATION.—SHADOW.—WHERE TO LOOK AND WHAT TO LOOK FOR.

Distance of Surgeon from Patient.—There is no fixed rule for this, and each surgeon may select his own distance. It might be well for the beginner to try different distances and then choose for himself. The writer prefers a one meter distance, and with few exceptions adheres to it. Some prefer six meters, others two meters, etc. The distance of one meter has important advantages: There is no getting up or down to place lenses in front of the patient's eye, as the patient or surgeon, or both, may lean forward for this purpose, if necessary. Another advantage is that at one meter distance there is a uniform allowance of one diopter in the estimate, which will be explained more fully under Rules for Retinoscopy at One Meter. To get the patient's eye and the observer's forehead just one meter apart, the distance may be marked off on the wall of the dark room on the side where the light is secured (see Fig. 7), or a meter stick for the purpose may be held in the hand of the observer or his assistant.

The method of obtaining the point of reversal



FIG. 6.—SHOWING DISTANCE FROM PATIENT'S EYES IN INCHES AND THE EQUIVALENT IN DIOPTERS.

at points other than the regulation one meter requires such an amount of extra measuring and computing that it does not meet with the general favor and satisfaction accorded to that found by producing an artificial myopia of one diopter. This can best be explained by reference to figure 6, where, if the observer is at one meter distance, and the neutralizing lens in front of the patient's eye focuses the emergent rays *about* that distance, he may have the liberty of moving forward five inches or backward five inches (a play of ten inches) in looking for the point of reversal, and not make a possible error in his result of more than twelve one-hundredths (0.12) of a diopter; *whereas* if he was working closer than this, say at half a meter, and was moved forward or backward five inches to find the point of reversal, he would likely make an error of 0.5 D., or even more, if he was not extremely careful in measuring the distance at which he found the reversal point.

Arrangement of Patient, Light, and Observer.—This has

already been described in great part, but reference to the accompanying sketch may give the student a more exact appreciation of the arrangement than any lengthy description could do.

For the convenience of the beginner in using the mirror, it is best, as here shown, to keep the surgeon's eye, the light, and the patient's eye on a hori-

FIG. 7.—ARRANGEMENT OF PATIENT, LIGHT, AND OBSERVER.

zontal line, and to accomplish this in children they will either have to stand, sit on a high stool, or on the parent's lap. The beginner will find it sufficiently difficult at first to reflect and keep the light on the patient's eye with the mirror held perpendicularly, without inclining it up or down, as he would have to do if the arrangement suggested is not carried out. Placing the light to one side of the

patient's head, or above it, and the observer seated at one meter distance from the patient, is a convenient way of working retinoscopy. It has two advantages: the observer avoids the heat of the flame, and at the same time does not have to move the light. But the writer is not partial to this mode of procedure, for various reasons of precision, explained in the text. Figure 8 shows the observer's eye, one meter from the patient's eye, and the light above.

FIG. 8.—LIGHT OVER PATIENT'S HEAD AND THE OBSERVER WITH MIRROR AT ONE METER DISTANCE.

Reflection from the Mirror.—The rays of light coming from the round opening in the screen to the plane mirror are reflected divergently, as if they came from the opening in the screen situated just as far back in the mirror as they originally started from in front (see Figs. 21, 23, and 25), and the patient, looking into the mirror, sees a round, bright spot of light, corresponding to the opening in the screen.

How to Use the Mirror.—It should be held

firmly before the right eye so that the sight-hole is opposite to the observer's pupil; and that it may be steady, the second phalanx of the thumb should rest on the cheek just below the eye, the edge of the metal disc even touching the side of the nose if the observer's interpupillary distance is not too great.



FIG. 9. FIG. 10.

AUTHOR'S MIRROR WITH FOLDING HANDLE.

FIG. 9.—Showing central light C, on small mirror B. This is the light the patient sees when looking into the mirror, and corresponds in size to the one-centimeter opening in screen. D is the folding cap handle to protect B when not in use. A is the metal disc.

FIG. 10.—Shows the light moved to one side as a result of tilting the mirror.

Thus held in position, its movements should be very limited, though they may be slow or quick, but *never*, at any time, should it be tilted more than one, two, or even three millimeters; for if inclined *more* than this the light is lost from the patient's eye. If the light,

the patient's, and the observer's eyes are on a horizontal line, then to find the patient's eye with the reflected light all the observer has to do is to reflect the light back into the light-screen, and by rotating the mirror to his right, carry the reflected light around on the same horizontal line until the patient's eye is reached. This may seem like a superabundance of instruction, but the finding of the patient's eye, which appears so easy, is an immense stumbling-block, at the beginning, to most students. Another way to find the eye is for the observer to hold his left hand up between his and the patient's eye and reflect the light on to it, and when this is done to drop his hand and let the light pass into the observed eye. Having succeeded in finding the patient's eye, the observer, if he is not very careful in his limited movements of the mirror and himself, will turn the light from the eye almost before he knows it, and so be compelled to start and find it again ; this causes much loss of time. A little practice on the schematic eye will assist the beginner wonderfully and give him courage, for if he hastens to the human eye, and then has to stop every minute or so to try and get the light on the eye, he soon becomes discouraged and shows his want of experience to the patient.

What the observer sees; or the general appearance of the reflection from the eye.—With the mirror held before his eye, and close up to the bright light coming from the ten-millimeter opening in the light-screen, the observer will obtain a reflection from the pupillary area of the patient's eye which varies in different patients, and is subject to certain changes

in the same patient as the refraction is altered by correcting lenses, or it may be changed by the turning of the patient's eye, or lengthening the distance between the mirror and the light, or increasing or diminishing the strength of the light, or increasing the distance between the observer and the patient. The reflection from the eye of the albino or blond is much brighter than from the brunette or mulatto, in whom it is not so bright, even dim. This character of the reflex is controlled, of course, in great part by the amount of pigment in the eye ground; however, in most instances there is more or less of an orange-red color to the reflex, and this is especially so as the point of reversal is approached; at the point of reversal, however, the reflex becomes less brilliant and possesses something of the color of a piece of newly coined silver. Cases of high errors of refraction give a dull reflex (see Fig. 19) as compared to low errors, where the reflex is usually *very* bright (see Fig. 11). Should the media be irregular or not perfectly clear, the reflex is altered accordingly; this will be referred to under the head of Irregular Astigmatism. The observer will also notice on the cornea and lens bright pin-point catoptric images, and at the inner edge of the iris, in many eyes, a very bright ring of light (see Fig. 11) about one millimeter in width, which is due to the very strong peripheral refraction; and as the eye is being refracted and the point of reversal approached, this peripheral ring may develop into a broader ring of aberration rays, which at times will be annoying. This will be referred to under Spheric Aberration, chapter VI.

Retinal Illumination.—By holding a strong convex lens closer to or further from a plane surface than its principal focus, or at the distance of its principal focus, and letting the sun's rays pass through it, there will be seen on the plane surface a round area of light; it is this light area which corresponds to the illumination on the retina, seen in retinoscopy by reflecting the light from the mirror into the patient's eye, and hence it is spoken of as the retinal illumination, the "illuminated area," "the area of light," "the image," etc.

FIG. 11.



FIG. 12.

FIG. 11.—UNIFORM ILLUMINATION IN AN EMMETROPIC EYE WITH SLIGHT SPHERIC ABERRATION.

FIG. 12.—UNIFORM ILLUMINATION AS IN FIG. 10, PASSED TO THE LEFT BY ROTATING THE MIRROR, DARKNESS OR SHADOW FOLLOWING.

Of course, the form of this illumination is controlled, in great part, by the refraction of the patient's eye.

Shadow.—This is the non-illuminated portion of the retina immediately surrounding the illumination. The retinal illumination and shadow are, therefore, in contact, and the contrast is most marked and easily recognized when the refractive error is a moderately high one, two or three diopters. It is by this combination of the illumination and non-illumination (shadow) that we study and give the "shadow test"

its name. In the dark room, the patient keeping his eye fixed, the retina is stationary and in total darkness, except the portion illuminated by the light from the mirror (see Fig. 11). If the mirror be rotated, the retinal illumination changes its place (see Fig. 12) and darkness, or shadow, appears in its stead. It is by this change of shadow (darkness) for illumination that we often speak of a movement of the shadow.

Where to Look and What to Look For.— With the patient, the observer, and the source of light in position as directed, the rays of light are reflected into the eye from the mirror as it is gently rotated in the various meridians, and the (1) *form*, (2) *direction, and* (3) *rate of movement* of the retinal illumination are carefully noted through a four- or five-millimeter area *at the apex of the cornea*, as *this* is the part of the refractive media in the normal eye that the patient will use *when* the effects of the cycloplegic pass away and the pupil regains its normal size.

The one- or two-millimeter area at the edge of the pupil should be avoided by the beginner, except in special instances, as only too frequently it contains a bright ring of light which may or may not give a stronger refraction than the area at the apex of the cornea (see Spheric Aberration, chap. VI).

The beginner will do good work with the retinoscope if he observes closely the illumination at the center of the pupil and *avoids* looking for shadows.

CHAPTER IV.

POINT OF REVERSAL.—TO FIND THE POINT OF REVERSAL.—WHAT TO AVOID.—DIRECTION OF MOVEMENT OF RETINAL ILLUMINATION.—RATE OF MOVEMENT AND FORM OF ILLUMINATION.—RULES FOR LENSES.—MOVEMENT OF MIRROR AND APPARATUS.

Point of Reversal.—This may be defined in several ways—namely: It is the far-point of a myopic eye, or

The artificial focal point of the emergent rays of light (Fig. 22), or

The point where the emergent rays cease to converge and commence to diverge, or

The point conjugate to a point on the retina (Fig. 26), or

The point where the erect image ceases and the inverted image begins, or

The point distant from the eye under examination, where the retinal illumination can not be seen to move.

The point of magnification.

To Find the Point of Reversal.—The recognition of the point of reversal is *the principle* of retinoscopy. It is what is sought for, and, when obtained under certain definite arrangements, is the correct solution of the test. During the test it is easy to tell when the illumination moves with or opposite to the movement of the light on the face, but to get the

exact point where there is no apparent movement is not always easy, and the ability to quickly find this point of reversal is only acquired after careful practice.

For example, having determined at one meter that the retinal illumination with a + 1.50 D. in front of the observed eye just moves with the light on the face, and against with a + 1.75 D., we know that the reversal point must be obtained with the lens numbered between the strength of these two lenses, *i. e.*, + 1.62 D. This demonstrates how we arrive at the exact correction, and also the capability and accuracy of retinoscopy.

Emmetropic and hyperopic eyes, in a state of rest, emit parallel and divergent rays, respectively, and to give such eyes a point of reversal, or a focus for the emergent rays, it will be necessary to intercept these rays with a convex lens as they leave the eye. In other words, emmetropic and hyperopic eyes must be made (artificially) myopic. In myopic eyes, however, the emergent rays always focus at some point inside of infinity, and the observer may, therefore, if he is so disposed, by moving his light and mirror to or from the patient's eye, as the case may be, find a point where the retinal illumination ceases to move. If this should be at two meters, the patient would have a myopia of 0.50 D.; if at four meters, a myopia of 0.25 D.; if at one meter, a myopia of one diopter, etc.

It is well for the beginner to remember, when using the plane mirror, that the illumination *on the patient's face always moves in the same direction the mirror*

is tilted, but *not* necessarily so in the pupillary area, where it may appear to move opposite; and *here* it is that we speak of the retinal illumination moving with or against (opposite to) the movement of the mirror, as the case may be, and make our diagnosis accordingly.

As the rays of light from the mirror proceed divergently to the patient's eye, as if they came from a point back in the mirror equal to the distance of the light (opening in light-screen) in front of it and working at one meter's distance, with source of light five inches in front of the mirror, the rays appear to emerge from a point five inches back of the mirror, or a total distance of 45 inches from the patient's eye, thus giving the rays of light a divergence equal to 0.87 of a diopter before they reach the patient's eye, and this point may be made conjugate to the retina. The observer will do good work if he reduces the retinal illumination to the utmost limit where it can be faintly seen moving with the movement of the mirror, and *if* this is done, the observer's eye and mirror will be just inside of the point of reversal where the erect image can still be recognized. In doing this, however, he must allow 0.87 in his estimate and not 1.0 D.

At the point of reversal no definite movement of the retinal illumination is made out and the pupillary area is seen to be uniformly illuminated, but not so brilliantly as when within or beyond the point of reversal.

If the observer's eye is, at this point, exactly conjugate to the retina, then the movement of the re-

flected light on the retina can not be perceived (though it does move), and the retinal illumination will occupy the entire pupil and the shadow will be absent.

Instead, however, of reducing the retinal illumination to the utmost limit (as just mentioned), where it can be faintly seen moving with the movement of the mirror, the writer prefers and recommends placing before the eye under examination such a lens or series of lenses as will bring the emergent rays of light to a focus on his own retina, so that no movement of the retinal illumination can be recognized.

When the point of reversal is approached, the uniform color of the retinal illumination occupies so much of the pupillary area that the student may think he has reached the point of reversal, and if he is not careful to pass the retinal illumination slowly across the pupil and get the shadow, he will find his result deficient, and possibly miss seeing some small amount of astigmatism.

To make sure that the point of reversal has been obtained, it is always best, especially for the beginner, to keep putting on stronger neutralizing lenses until he gets a *reversal* of movement, when he knows at once that the point of focus of the emergent rays has passed in between the mirror and eye under examination.

The lenses which control the rays of light emerging from the patient's eye are spoken of as neutralizing lenses.

What to Avoid.—It occasionally happens that a retinal vessel or vessels or a remnant of a hyaloid

artery, if present, or even the nerve head, may be seen when the light is reflected into the eye; if so, they are to be ignored, as they are not parts of the test. If the patient's eye is turned, or the rays from the mirror fall obliquely, or the neutralizing lens in front of the eye is inclined instead of being perpendicular, there will be seen reflections of light and images upon the neutralizing lens or cornea, or both, and, in consequence, the retinal illumination is more or less hidden or obscured; these images and reflections can be easily corrected by removing the cause. The catoptric images can not be removed, but as they are very small, the beginner soon learns to ignore them. The retinal illumination may occasionally contain a small dark center, which will disturb the beginner unless he remembers that it is caused by the sight-hole in the mirror, and is likely to occur when the sight-hole is large and cut through the mirror. This same dark center in the illumination is also seen at times when the light is removed some distance from the mirror, and the correcting lens almost neutralizes the refraction. The neutralizing lens should never be so close to the eye that the lashes touch, and, in warm weather especially, moisture from the patient's face may condense on the trial-lens, and temporarily, until it is removed, obscure the reflex.

Retinoscopy with a Plane Mirror at One Meter's Distance and Source of Light close to the Mirror.—Direction of Movement of Retinal Illumination.—Rate of Movement and Form of Illumination.—These important points in refer-

ence to the retinal illumination should be decided promptly and without any prolonged examination. This proficiency, of course, will only come by practice, and if, on first examination, the observer can not decide whether the retinal illumination is moving with or opposite to the movement of the reflected light on the face, he may approach the eye until this point is determined. At the distance of one meter the three important essentials may be stated in the following order and in the form of rules:

Direction of Movement of Retinal Illumination.—The recognition of the direction that the retinal illumination takes when tilting the mirror is a most important point in the study of retinoscopy.

The movement of the retinal illumination, when rotating the mirror, going *with* the movement of the light on the patient's face, signifies emmetropia, hyperopia, or myopia, if the myopia be *less* than one diopter.

The apparent movement of the retinal illumination going *opposite* to the movement of the light on the face always signifies myopia of *more* than one diopter.

Rate of Movement.—This, of course, is under the control and is influenced in great part by the rate of movement of the mirror itself; yet after a little practice the observer will recognize the fact that there is a certain slowness in the apparent rate of movement of the illumination when the refractive error is a high one and requires a strong lens for its neutralization, whereas when the retinal illumination appears to move fast, the refractive error is

but slight, and requires a weak lens for its correction.

Form of Illumination.—A large, round illumination, while it *may* signify hyperopia or myopia alone, yet it does not preclude astigmatism in combination.

When the illumination appears to move *faster* in one meridian than the meridian at right angles to it, astigmatism will be in the meridian of *slow* movement. If the retinal illumination is a band of light extending across the pupil, it signifies astigmatism.

The width of the band of light does not indicate



FIG. 13.—STRAIGHT EDGE, INDICATING ASTIGMATISM.



FIG. 14.—CRESCENT EDGE, INDICATING SPHERIC CORRECTION.

so much the strength of the correcting cylinder required for its neutralization as does the apparent rate of movement; if slow, a strong, if fast, a weak, cylinder is required.

The band of light that is seen when a spheric lens of one diopter or more corrects one meridian, and the meridian at right angles remains partly corrected, indicates the axis of the cylinder in the prescription.

Rules for Placing Neutralizing Lenses.—A *plus* lens is required when the retinal illumination moves *with* the illumination on the face, and a *minus*

lens is required when it moves *opposite* to the light on the face.

Movement of the Mirror.—There are times when a quick movement, and, at other times, a slow or gradual movement, is required. A substitution of the quick for the slow movement, and the result can not always be accurately determined. This is explained under "slow movement."

FIG. 15. — WÜRDEMANN'S DISC.

A quick movement of the mirror may be used when looking into the eye before any correcting lens has been placed *in situ*. It often tells the character of the refraction.

The slow movement of the mirror and the five-millimeter opening in light-screen come into use and are of the *utmost* importance when the eye has been corrected to within 0.75 D. or less, as it is generally at this point that so many, by a *quick* movement, hasten the peripheral rays and mask the central illumination, giving the idea at once of over-correction (see Spheric Aberration, chap. VI). This is a most common error with the beginner, the inexperienced, and with those who fail to get good results and who ridicule retinoscopy as "not exact," or as "not agreeing with the subjective method." It is well in *every* instance, when the point of reversal is approached, to pass the retinal illumination (not

the light area on the face) well across the pupillary area to make *sure* in regard to the character of shadow which follows or precedes it. This movement, at such a point in neutralization, will often give a hint as to the presence of astigmatism or not, as a reference to figures 13 and 14 will show.



FIG. 16.—JENNINGS' SKIASCOPIC DISC.

The presence of astigmatism is known by the straight edge of the illumination, or, in its place, a crescent edge would mean a spheric correction.

Apparatus for placing lenses in front of the patient's eye.—There are several different forms in the market, their purpose being twofold—to save time and any extra movements on the part of the

surgeon. Of these, that of Würdemann (*American Journal of Ophthalmology*, p. 223, 1891) seems the best hand skiascope. A reference to the sketch (Fig. 15) shows this instrument with its convenient handle, whereby the patient, being instructed, raises or lowers the disc in front of the eye, with its smooth broad edge resting against the side of the nose.

One column contains plus and the other minus lenses, and as it is reversible, these may be placed in front of the eye, as the surgeon directs.

The most modern and complete revolving skiascopic disc is that of Jennings (*American Journal of Ophthalmology*, November, 1896, and April, 1899), and may be best understood from his own description: "It consists of an upright metal frame, 18 inches high and 7 inches wide, placed at the end of a table 26½ inches long and 12 inches wide. In the upright frame is an endless groove containing 39 lenses and 1 open cell. At the lower end of the frame is a strong driving wheel connected with a horizontal rod running the length of the table to a handle with which the operator rotates the lenses. Facing the operator and close to his hand is a large disc, on which is indicated the strength of the lens presenting at the sight-hole. The white numbers on a black ground represent convex, and the black numbers on the white ground concave, lenses. The lenses range from 0.25 D. to 9 D. plus, and from 0.25 D. to 9 D. minus. The sight-holes are ⅞ of an inch in diameter, and are placed about five inches from the top of the upright frame. In

FEB 15 1923

front of each sight-hole is a cell marked in degrees to hold stronger spheres or cylinders. The central portion of the upright is cut away, leaving a space for the face of the patient. A movable blinder is hung from the top, while the chin-rest moves up and down on two parallel rods and is held in place by a thumb-screw. The whole is mounted on a strong adjustable stand, which is raised or lowered by means of a rack and pinion. The essential advantages of this skiascope are as follows:



FIG. 17.—AUTHOR'S TRIAL FRAME WITH AXONOMETERS ATTACHED.—(*Drawing reduced in size.*)

1. It saves time and fatigue in changing lenses.

2. It is *under the immediate control of the operator*, and indicates the lens in front of the sight-hole without his getting up.

3. The mechanism is simple, durable, and easy to operate.

4. The cornea is accurately centered and the lens perpendicular to the front of the eye (a very important consideration, and one not possible with *every* kind of trial frame).

5. The instrument is of such length that the

operator is always one meter distant from the patient.

While either the hand or the revolving disc is recommended, yet the writer is partial to an accurately fitting trial frame (Fig. 17), using the lenses from the trial-case, which should be conveniently at hand. The following suggestions in the selection and use of the trial frame are offered: The temples should rest easily on the ears, the nose-piece (bridge) to have a sufficiently long post to permit the eye-pieces to fit high and accurately over any pair of eyes, especially those of children, and have the corneæ occupy the center of each eye-piece. Correct results can not be expected or quickly obtained unless the neutralizing lenses be placed with their centers corresponding to corneal centers, and at the same time perpendicular to the front of the eye. A convenient and small trial-case containing a row of plus and minus spheres, from 0.12 to 10 D., is shown in figure 18.



FIG. 18.—AUTHOR'S TRIAL-CASE FOR RETINOSCOPY.

CHAPTER V.

RETINOSCOPY IN EMMETROPIA AND THE VARIOUS FORMS OF REGULAR AMETROPIA.—AXONOMETER.

Hyperopia.—In this form of refraction the direction of the movement of the retinal illumination is *with* the movement of the light on the patient's face. By rotating the mirror in the various meridians and observing the rate of movement, a strong or weak plus sphere, according to the apparent rate of move-

FIG. 19.

FIG. 20.

FIG. 19.—GRAY REFLEX AS SEEN IN HIGH HYPEROPIA, EVEN DARKER THAN THE PICTURE SHOWS IT.

FIG. 20.—GRAY REFLEX, WITH CRESCENT EDGE BY TILTING MIRROR TO LEFT, DARKNESS OR SHADOW FOLLOWING.

ment, is placed before the eye, and the rate of movement of the retinal illumination is again noted.

Practice alone will guide the observer in a quick appreciation of the approximate strength of neutralizing lens to thus employ.

If the movement of the illumination appears slow, and the observer places a + 2.75 D. before the eye for its neutralization, and the illumination then becomes brilliant and appears to move fast and

with the light on the face, the hyperopia is still slightly uncorrected and a stronger lens must be substituted. (At this point in the examination the five-millimeter opening in the light-screen may be used to advantage.)

Removing the + 2.75 D. and substituting a + 3.25 D., if the retinal illumination is found to move *opposite* to the movement of the light on the face, the refraction of the eye will then be between the + 2.75 D. and the 3.25 D., which is 3 D. (See example, p. 41, chap. IV.) Now, while the + 3 D. has brought the emergent rays to a focus at one



FIG. 21.

meter, it has made the eye myopic just one diopter, so that in taking the patient from the dark room to test his vision at six meters, or infinity, this one diopter (artificial myopia) must be subtracted from the + 3 D., which would leave + 2 D., the amount of the hyperopia.

A reference to figures 21 and 22 will illustrate the description just given.

Figure 21 is the hyperopic eye under examination, and shows the mirror at one meter's distance, with the light five inches from the mirror. The dotted lines represent the rays proceeding divergently

from the eye under examination ; the dark lines show the reflected rays from the mirror, which illuminate the retina and have an imaginary focus (dotted lines) beyond the retina.

Figure 22 is a profile view showing the hyperopic eye with neutralizing lens in position. The dotted lines with arrow-heads indicate the direction the rays would naturally take coming from the eye. The lens (+ 3 D.) in front of the eye is just sufficiently strong to bend these divergent rays and bring them to a focus at one meter's distance (artificial point of



FIG. 22.

reversal). In other words, + 2 D. of the three diopters thus placed before this hyperopic eye would have bent the divergent rays and made them parallel, or emmetropic, but the additional one diopter bends the rays still more and brings them to a focus (P. R.) at one meter. If, now, the observer approaches the eye thus refracted and observes the retinal illumination closer than one meter, he will be inside of the point of reversal, and consequently see an erect image moving rapidly with the direction of the movement of the mirror. If beyond this point of reversal, he would get an inverted

image and the retinal illumination moving rapidly in a direction opposite to the movement of the mirror.

Emmetropia.—The emergent rays from an emmetropic eye are always parallel, and the observer seated at one meter sees the pupillary area in such an eye brilliantly illuminated, the illumination moving rapidly with the light on the face as the mirror is slowly rotated.

A reference to figure 23 shows the emmetropic eye under examination with the position of light, mirror, and eye, as in figure 21. The dotted lines



FIG. 23.

indicate the parallel emergent rays, and the solid lines the divergent rays from the mirror with an imaginary focus just beyond the retina. The purpose in this instance, as in all others of retinoscopy, is to place such a neutralizing lens before the eye as will bend the emergent rays and bring them to a focus at a certain definite distance, making the emergent rays from a point on the retina come to a focus on the observer's retina. Therefore, to change this illumination so that no movement can be seen to take place in the pupillary area, and at the same time have the emergent rays focus on the

observer's retina, a + 1 sphere must be placed before the eye.

Just here the writer wishes to *impress* upon the beginner the great importance, as mentioned on page 30, of refracting the macular region. To accomplish this, the patient must fix his gaze upon the metal disc of the mirror. As the region of the macula is departed from, the strength of the neutralizing lens grows slightly stronger in emmetropia and hyperopia, and diminishes in myopia. A reference to figure 24 will give an idea of what is meant, and show that radii drawn from the nodal point are not the same length as the one to the fovea.



FIG. 24.

Myopia.—In myopia the emergent rays *always* converge to the far-point (point of reversal), and the observer, seated at one meter distant from the eye, will have the apparent movement of the retinal illumination going opposite to the light on the face *if* the myopia exceeds one diopter, and *with* the light on the face *if* the myopia is less than one diopter. If the myopia should be *just* one diopter, then the emergent rays would focus on the observer's retina at one meter, and there will not be any neutralizing lens required to accomplish this purpose; but if the emergent rays focus beyond one meter, the observer will be within this point of reversal or focus, and will, therefore, have an erect image, moving fast with the movement of the mirror, and will have to place before the eye a plus lens of less than one diopter to

bring the point of reversal up to the distance of one meter. When the myopia is more than one diopter, and observer at one meter, the emergent rays will have focused somewhere between the observer and the patient, and, as a result, the retinal illumination appears to move opposite to the light upon the face, more or less rapidly, according to the amount of myopia ; and a concave or minus lens must be placed in front of such an eye that will bring the emergent rays to a focus at one meter, or, in other words, will stop all apparent movement of the retinal illumina-



Fig. 25.

tion. If, for example, a —2.75 D. has been so placed, and the movement is still slightly opposite to the movement of the mirror, and a —3.25 D. substituted makes the retinal illumination move *with* the movement of the mirror, then the neutralizing lens for one meter will be between —2.75 D. and —3.25 D., which will be —3 D.

Figure 25 shows the myopic eye just described, with the position of the mirror, light, and eye, as in figures 21 and 23. The solid lines represent the rays reflected divergently from the mirror focusing at a point in the vitreous before coming to the retina, and

the broken lines show the rays emerging from a point on the retina and then converging to the focus, far-point, or point of reversal close to the eye, between the eye and the mirror. The observer, seated with the mirror one meter distant, gets an opposite movement in the pupillary area from the direction in which he moves his mirror, and, of course, an inverted image. If the observer had his eye at the point where the emergent rays focused (dotted lines cross), he would not recognize any movement in the pupillary area, and it would have a uniform reflex. The amount of the myopia is equal to the distance meas-



FIG. 26.

ured from this point of reversal to the cornea ; for example, if the distance (point of reversal) was twenty-five cm. from the patient's eye, then the amount of the myopia would be four diopters ; if at 33 cm., then 3 D., etc.

Figure 26 is a profile view of the myopic eye. The dotted lines show the rays coming from a point on the retina and focusing at the far-point (F.P.) ; the solid lines show the emergent rays acted upon or bent by a plano-concave lens of three diopters, which has lessened the convergence of these emergent rays and put the far-point farther from the eye,

or at a distance of one meter. The observer at this distance does not appreciate any movement in the pupillary area, but if he moves the light and mirror closer to the eye he is then inside the point of reversal and gets an erect image moving with the movement of the mirror; if beyond the one meter's distance, an inverted image and movement against the movement of the mirror will be seen. If a —4 D. lens had been placed before this myopic eye, the emergent rays would have proceeded from it parallel, and the observer, at one meter, would have the same conditions as in the refraction of an emmetropic eye, figure 23; but as only a —3 D. glass was used, the eye has one diopter of its myopia uncorrected. From the description of retinoscopy in hyperopia, emmetropia, and myopia, just given, the student will recognize at once that the hyperopic, emmetropic, and myopic eye of less than one diopter, working with the plane mirror at one meter's distance, are given a stronger refraction than they naturally call for, or, in other words, are made, artificially, myopic one diopter. And the myopic eye of more than one diopter, under similar conditions, being already myopic, retains one diopter of its myopia. To give a patient thus refracted with the retinoscope his emmetropic correction (correction for parallel rays of light), *an allowance must always be made, in all meridians, of one diopter, no matter what the refraction.* The artificial myopia thus produced at one meter gives the following rules for glasses required for infinity:

Rules.—1. When the neutralizing lens employed is plus, then subtract one diopter.

2. When the neutralizing lens employed is minus, then add a —1 D., or what is more simple, or even a better rule, is, To *always add a —1 sphere to the neutralizing lens obtained in the dark room when working at one meter, and the result will be the emmetropic or infinity correction.*

Examples:

Dark Room, ...	+0.50	0.00	+1.00	+2.00	—1.00
Adding, ...	—1.00	—1.00	—1.00	—1.00	—1.00
Emmetropic Correction,	—0.50	—1.00	—0.00	+1.00	—2.00

The main point in all retinoscopic work to remember in changing from the dark room to the six meter correction, is to always allow for the distance from the patient's eye to the point of reversal—i. e., if working at half a meter, allow two diopters; if at two meters, 0.50 D.; if at four meters, 0.25 D., etc.

Regular Astigmatism.—When refracting with the retinoscope, the observer should remember *that he is refracting the meridian in the direction of which he moves the mirror.* Particular attention is called to this important fact on account of the confusion sometimes arising in the student's mind from the use of the ophthalmoscope, where the refractive condition of a certain meridian is estimated by the strength of the lens used to see clearly the vessels at right angles to it. Astigmatism being present in an eye, means a difference in the strength of the glass required for the two principal meridians, which, with few exceptions, are at right angles to each other, and it is to these two principal meridians *only* that the

observer pays attention; for example, the eye that take the following formula,

$$+ 1.00 \text{ D.} \bigcirc + 1.00 \text{ c. axis } 105°,$$

means that in the 105 meridian there is + 1 D. and in the 15 meridian a + 2 D. In the dark room a + 2 sphere in front of such an eye at one meter would correct the 105 meridian and partly correct the 15 meridian; or a + 3 D. would correct the 15 and over-correct (movement against) the 105 meridian. When with + 2 D. the 105 meridian is cor-



FIG. 27.

rected and the 15 only partly so, there is seen in the 15 meridian a band of light which stands or extends across the pupil in the 105 meridian and moves across the pupil from left to right *with* the movement of the mirror as it is tilted in the 15 meridian.

The presence of this band of light *after* the meridian of least ametropia has been corrected *always* signifies astigmatism, and the axis it subtends—in this case 105°—gives the axis of the cylinder in the prescription; and the amount of the astigmatism, or the strength of the cylinder required, is the differ-

ence between the strength of the two spheres employed. Figure 27 shows the method of writing such a dark room correction, and adding, according to our rule, a —1 to this dark room work, we get our original formula:

$$+ 1.00 \text{ D. } \bigcirc + 1.00 \text{ c. axis } 105°.$$

The method of correcting with spheres (Fig. 18) will be found much more satisfactory than by placing a + 2 D., as called for in the 105 meridian, then adding and changing cylinders until the correct one is found.

FIG. 28.

FIG. 29.

FIG. 28.—BAND OF LIGHT AT AXIS 60°, WITH THE 60° MERIDIAN NEUTRALIZED. No movement of the illumination can be recognized in the 60° meridian.

FIG. 29.—Shows the same as figure 28, but the band of light with straight edge has been moved upward and to the left by tilting the mirror in the 150° meridian.

It takes much time and care to get the cylinder axis just right, and is most difficult in the dark room. After the result has been obtained with spheres, the observer may, if he is so disposed, prove it before leaving the dark room with the sphero-cylinder combination.

Astigmatism may or may not be recognized on first inspection of the fundus-reflex, this depending *entirely* on the refraction; if it be a high astigmatism with a small amount of refractive error in the op-

posite meridian, as in one of the following formulas,

$$+ 1.00 \text{ D. } \bigcirc + 3.00 \text{ c. axis } 45°,$$
$$- 1.00 \text{ D. } \bigcirc - 4.00 \text{ c. axis } 180°,$$

then the band of light so characteristic of astigmatism will be plainly seen on first inspection, extending across the pupil before any neutralizing lens has been placed in position; but if the hyperopia or myopia be high and the cylinder required is low, as in one of the following formulas,

$$+ 3.00 \text{ D. } \bigcirc + 0.75 \text{ c. axis } 105°,$$
$$- 4.00 \text{ D. } \bigcirc - 1.00 \text{ c. axis } 165°,$$

then the band of light is not recognized on first inspection or until an approximate correction has been placed before the eye. To get an idea of what the band of light looks like, the beginner may refer to figures 28 and 30; or focus rays of light through a strong cylinder; or place a cylinder in front of the schematic eye and study the retinal illumination. The student should bear in mind that the axis of the band of light appears on the meridian of least ametropia, and is brightest when this meridian has received its full spheric correction—the opposite meridian being only partly corrected.



FIG. 30.—BAND OF LIGHT. ASTIGMATISM AXIS 90°.

The reason for the brightness of the band of light when the meridian of its axis is corrected is that any point on the retina in this meridian is conjugate to the focus on the observer's retina (point of re-

versal), and any movement of the mirror in this meridian is not recognized, but has a uniform color and occupies the entire meridian of the pupil. To recognize so small an error as a quarter diopter cylinder,—which is *not* easily detected, and the observer, if he is in a hurry, might think the case one of simple hyperopia or myopia,—the writer would suggest that when the supposed point of reversal is reached the correcting sphere be increased a quarter of a diopter, and *if* only one meridian is found over-corrected (movement opposite), the other remaining correct (no movement

FIG. 31.—BAND OF LIGHT SHOWING HALF A DIOPTER OF ASTIGMATISM.

recognized), he *then* knows that a quarter cylinder is required; for example, a + 2 D. is supposed to correct all meridians, and yet by substituting a + 2.25 D., the vertical meridian moves against and the horizontal remains stationary; then a + 0.25 D. cylinder is called for at axis 90°.

Cases having a low astigmatic error of 0.50 D. can be recognized when near the point of reversal by the faint shaded area on each side of the band of light, as shown in figure 31—a condition often overlooked.

Mixed Astigmatism.—In this condition of refraction, where one meridian is myopic and the

meridian at right angles to it is hyperopic, the movement of the retinal illumination in the myopic meridian will be controlled by the *amount* of the myopia. The illumination in the myopic meridian, if the myopia is less than one diopter, moves *with* the mirror, and against the movement of the mirror if it is more than one diopter; in either instance the observer gets a distinct band of light in the meridians alternately as each meridian is neutralized separately with a sphere. Taking the following example,

—2.00 c. axis 180° ⊖ + 1.00 c. axis 90°,

the 90 meridian shows an *opposite* movement, and in the horizontal the movement is *with* the movement of the mirror. If, now, a —1 D. sphere be placed before the eye, the 90 meridian is neutralized for one meter distance, and a bright band of light is seen at 90°, moving with the movement of the mirror in the horizontal meridian. Removing the —1 D. and placing a + 2 D. before the eye, which would neutralize the horizontal meridian for one meter, a bright band will be seen in the horizontal meridian and moving opposite to the movement of the mirror in the 90° meridian. Carrying out the rule of always adding a —1 D. sphere to the correction obtained in the dark room at one meter, we have —1 added to the —1 in the vertical meridian, making —2 D., in axis 180°; and adding —1 to the + 2 D. in the horizontal, we have + 1 D. in axis 90°, or our original formula:

—2.00 axis 180° ⊖ + 1.00 c. axis 90°.

The rule for neutralizing lenses in mixed astigma-

tism is the same as for any other form of refraction; namely, using a plus lens when the movement is with, and a minus lens when the movement is opposite to, the movement of the light on the face.

To transpose crossed cylinders into a sphero-cylinder combination the writer would advise using the rule of Dr. Harry S. Pearse, of Albany, which is as follows:

"The cylinder is the sum of the two cylinders with the sign and axis of one of the cylinders. The sphere is the strength of the other cylinder with its sign." In the above formula, the sphero-cylinder combination will be

$$-2 \text{ D. } \bigcirc +3.00 \text{ c. axis } 90°.$$

Axonometer.—To find the exact axis subtended by the band of light while studying the retinal illumination, when the meridian of least ametropia has been corrected, the writer has suggested a small instrument which, for want of a better name, he has called an axonometer.

Figure 32 shows this instrument, and figure 33 the axonometer in position.

The original description of this device was published in *The Medical News*, March 3, 1894, as follows: "The direction of the principal meridians of corneal curvature is often difficult to determine, and the statement of the patient must be accepted when confirming the shadow-test correction; or, if there is still uncertainty, the ophthalmometer of Javal may be of service. The axonometer is a black metal disc, with a milled edge, one and one-half mm. in thick-

ness, of the diameter of the ordinary trial-lens, and mounted in a cell of the trial-set. It has a central

FIG. 32.

round opening 12 mm. in diameter—the diameter of the average cornea at its base. Two heavy white lines, one on each side, pass from the circumference across to the central opening, bisecting the



FIG. 33.

disc. To use the axonometer, place it in the front opening of the trial frame, and with the patient

seated erect and frame accurately adjusted so that the cornea of the eye to be refracted occupies the central opening, proceed as in the usual method of making the shadow test. As soon as that lens is found which corrects the meridian of least ametropia, and the band of light appears distinct, turn the axonometer slowly until the two heavy white lines accurately coincide, or appear to make one continuous line with the band of light (see Fig. 33).

"The degree marks on the trial frame to which the arrow-head at the end of the white lines then points is the exact axis for the cylinder. The axonometer possesses the following points of merit:

"Simplicity.

"Accuracy.

"Small expense.

"It covers an unnecessary part of the trial-lens which too frequently gives annoying reflexes and images.

"It saves time, avoids the statement of the patient, and renders the ophthalmometer unnecessary.

"Its color (black) absorbs the superfluous light rays from the mirror and gives a stronger contrast to the reflex and central illumination.

"Limiting the field of vision in children, it permits of more concentrated attention.

"For children and nervous patients, when it is difficult to use the ophthalmometer, this simple appliance is of great service."

Lately the writer has improved the axonometer by having the white lines broadened to four millimeters, which is a decided advantage over the

instrument shown in figures 32 and 33, as the broad line is easily seen at one meter distance. This

FIG. 34.

axonometer, shown in figures 17 and 34, is made of thick celluloid.

CHAPTER VI.

RETINOSCOPY IN THE VARIOUS FORMS OF IRREGULAR AMETROPIA. — RETINOSCOPY WITHOUT A CYCLOPLEGIC. — THE CONCAVE MIRROR. — DESCRIPTION OF THE AUTHOR'S SCHEMATIC EYE AND LIGHT-SCREEN.—LENSES FOR THE STUDY OF THE SCISSOR MOVEMENT, CONIC CORNEA, AND SPHERIC ABERRATION.

Irregular Astigmatism.—This condition is either in the cornea or in the lens, or in both structures in one and the same eye; in any instance it is confusing to the beginner, and even the expert must work slowly to obtain a definite result. The corneal form is most difficult to refract, as the retinal illumination is more or less obscured by areas of darkness. The illumination between these dark areas appears to move with, in places, and in others against, the movement of the mirror. By moving the mirror so as to make the light describe a circle around the pupillary edge, a most unique kaleidoscopic picture is obtained, which is quite diagnostic of the condition. To refract an eye with this irregularity the observer may have to change his position several times, going closer to or farther away from the patient. Very often these eyes are astigmatic, and the band of light may be promptly noted by the observer changing his position as suggested, and at the same time placing a neutralizing lens in position. Care *must* be taken, also, to refract in the area of the cornea that

will correspond to the small pupil when the effect of the cycloplegic passes away. It is often best, in these cases of irregular corneal astigmatism, to make a record of the correction found and use it as a guide in a post-cycloplegic manifest refraction.

Irregular astigmatism of the lens is frequently more or less uniform, and not so broken as in the corneal variety. Figures 35 and 36 show two kinds of irregular lenticular astigmatism.

Figure 35 illustrates the spicules pointing in from the periphery, and so long as these do not encroach upon the pupillary area, they do not usually in them-

FIG. 35.

FIG. 36.

IRREGULAR LENTICULAR ASTIGMATISM.

selves interfere with vision; they are not often recognized until the pupil is dilated, are then very faint, and not usually made out until the point of reversal is approached. Figure 36 is another form of irregular lenticular astigmatism, and a very interesting picture as studied with the retinoscope; and, as in figure 35, *when* very faint, is not made out until close to the point of reversal. These two forms of irregular lenticular astigmatism, when *just* beginning, are very seldom seen with the ophthalmoscope; the striations are too fine to be made out except under the conditions just described, and when recognized are of inestimable value from a point of

prophylactic treatment, calling for a change of occupation, rest to the eyes, and carefully selected glasses, the latter often being weak lenses. These lenticular conditions not infrequently accompany the "flannel-red" fundus, the "fluffy eye ground," the "shot-silk retina," the "woolly choroid," etc.

Scissor Movement.—Another form of astigmatism that may be classed as irregular is where there are two areas of light, each with a straight edge, and usually seen on the horizontal meridian, or inclined a few degrees therefrom either way, and



FIG. 38.—LIGHT AREAS COMING TOGETHER AND DARK INTERSPACE FADING.

moving toward each other as the mirror is tilted in the opposite meridian; in other words, as the observer is seated at one meter he sees an area of light above and an area of light below with a dark interspace (Fig. 37). As the mirror is slowly tilted in the vertical meridian these light areas approach and are followed by darkness or shadow, and at the same time the dark interspace begins to fade, giving the picture as shown in figure 38. When the light areas are brought together, they result in a horizontal band of light, as seen in figure 39, and at this point resemble the ordinary band of light as seen in

regular astigmatism. This movement of the light areas is likened to the opening and closing of the scissor blades, and hence the name of scissor movement.

These cases are more or less difficult to refract, but the presence of the two areas of light with the dark interspace will often assist in a correct selection of glasses, for while they are generally of the compound hyperopic variety, calling for a plus sphere and plus cylinder, yet practice and the patient's

FIG. 37.

FIG. 39.

FIG. 37.—LIGHT AREA ABOVE AND BELOW, WITH DARK INTERSPACE.
FIG. 39.—LIGHT AREAS BROUGHT TOGETHER.

statement often call for a plus sphere and minus cylinder.

With the following formula,

$$+ 2.00 \text{ D. } \bigcirc + 0.75 \text{ c. axis } 90°,$$

substituting a sphere the strength of the combined values of the sphere and cylinder, and using a minus cylinder of the same number as the plus cylinder at the opposite axis, the result will be,

$$+ 2.75 \text{ D. } \bigcirc - 0.75 \text{ c. axis } 180°.$$

The vision with the latter formula is much better in many instances than with the former, and though either formula would be correct, yet the latter is practically the better of the two, and should be ordered when so found. The condition which may

be the probable cause of the scissor movement is a slight tilting of the lens (see Fig. 40)—that is, the antero-posterior axis of the lens does not stand perpendicular to the plane of the cornea, thus making one portion of the pupil myopic (area of light moving opposite) and the other portion hyperopic (area of light moving with the movement of the mirror). This condition may be simulated by placing a convex lens at an angle before the schematic eye, or reflecting the light into the eye obliquely, or by using the combination lens in front of the schematic eye, as suggested on page 83. What



FIG. 40.

causes the tilting of the lens the writer is not prepared to state *positively;* it may be congenital, and yet careful inquiry of the patients, in many instances, has shown that it is most likely due to using the eyes to excess in the recumbent posture. It may be a coincidence, but most of the cases of scissor movement seen by the author have been in adults, and those who were in the habit of reading while lying down, reading themselves to sleep at night in bed.* Other cases were seen among

* The writer does not wish to be misunderstood and does *not* say that every one who uses his eyes in this faulty position *must* develop this form of irregular astigmatism.

paper-hangers, whose occupation compelled them to look upward much of the time. These do not seem unlikely causes, especially when the anatomy of the ciliary region is considered, the strain of the accommodation (possibly spasm) during the faulty position of the eye tilting the lens as it rests upon the vitreous body. This form of astigmatism, so far as known, remains a permanent one even after a cessation from the original cause and correcting glasses have been ordered. The retinoscope is the only instrument of precision we have in diagnosing this condition. The ophthalmoscope may recognize



FIG. 41.

the presence of the astigmatism, but not its character, and the ophthalmometer only records the corneal curvature. Cases of aphakia (following cataract *extraction*) frequently show the scissor movement *during* the process of retinoscopy. This is undoubtedly due to the flattening of the cornea corresponding to the section, making one portion myopic and the other hyperopic. Figure 41, with correcting sphere in position, shows such a condition, where the upper illumination would move with and the lower, being myopic, would move against the movement of the mirror.

Compound Irregular Astigmatism.—This is a

combination of the scissor movement and regular astigmatism, but they are *not* at right angles to each other. The scissor movement may be at 180°, and the regular astigmatism at some point *away* from 90°, but not at 90°; or the regular astigmatism may be at 90° and the scissor movement at some meridian other than 180°.

A hasty review of the literature of astigmatism does not reveal any reference to this form, and the name for the condition has been suggested by the following picture, namely: When studying the reflex, a vertical band of light will be seen passing across the pupillary area from left to right as the mirror is turned, and then in the vertical meridian (*not* at right angles) the scissor movement will be recognized also; there is, therefore, a combination of regular corneal astigmatism with the scissor movement at an oblique angle, giving the compound name suggested. This form of astigmatism is rare, yet not difficult to diagnose or refract when understood. It is hoped, however, that the beginner in retinoscopy may not meet one of these on his first attempt at the human eye. (See page 85.)

Conic Cornea.—Reflecting the light into an eye that has such a condition, the observer is impressed at once with the bright central illumination that moves opposite to the movement of the mirror, the peripheral illumination moving with, unless perchance the margin should be myopic also, but of less degree. This form of illumination is seen in figure 42, showing the central illumination faintly separated by a shaded area or ring from the peripheral circle. The best

way to refract a case of this kind is to keep a record of the neutralizing lens or lenses required for the portion of the pupillary area that will correspond to the size of the pupil after the effect of the cycloplegic passes away, and use this record as a guide in a post-cycloplegic manifest correction, as in irregular corneal astigmatism.

As the apex of the cone is not always central, the observer must not expect to always find the bright illumination in the center of the pupillary area, as just mentioned; and it is also well to note the fact that a band of light will often appear during the

FIG. 42.—ILLUMINATION SEEN IN CONIC CORNEA.

process of neutralization, as astigmatism is usually present. This is further described on page 84.

Spheric Aberration.—This appears under two forms, positive or negative, and is the condition in which, during the process of neutralization, there are two zones, one central and the other peripheral, where the refraction is not the same. In positive aberration the peripheral refraction is stronger and in negative aberration the peripheral is weaker than the central area; that is to say, in the positive form, when the point of reversal for the center of the pupil is close to one meter, the peripheral illumination grows broader and has a tendency to, and often will,

crowd in upon the small central illumination, giving the idea of neutralization, or even the appearance of over-correction, the illumination in the periphery moving opposite. The observer must be on his guard for this condition, and while giving the mirror a slow and limited rotation must watch carefully the



FIG. 43.—POSITIVE ABERRATION.

illumination in the center of the pupil and *not* hasten the peripheral movement. (See What to Avoid, p. 43, chap. IV.) The observer may have to approach the patient's eye closer than one meter if the peripheral illumination appears to move very fast. The negative form is where the peripheral refraction is



FIG. 44.—NEGATIVE ABERRATION.

weak as compared to the central, which appears strong, and when the neutralizing lens gives a point of reversal at the center of the pupil the peripheral illumination still moves with the movement of the mirror. This condition is seen in cases of conic cornea.

Figure 43 illustrates positive aberration where the

parallel rays passing through a convex lens in the periphery at *A A* come to a focus at *A'*, much sooner than the parallel rays *B B*, near the center, which come to a focus back of *A'* at *B'*.

Figure 44 illustrates negative aberration, which is the reverse of positive aberration, and the central rays *B B* are focused at *B'* in front of the peripheral rays *A A* focusing at *A*.

Retinoscopy Without a Cycloplegic.—Cases of myopia and mixed astigmatism which have large pupils *can* be quickly and accurately refracted by the shadow test without the use of a cycloplegic. This has been repeatedly proven by comparison of the manifest and cycloplegic results; yet it is not a method to be recommended or pursued, for two reasons: One is that these patients are not annoyed, like hyperopics, by the blurred near-vision incident to the cycloplegic; and, secondly, glasses ordered without the cycloplegic seldom give the comfort that follows from the physiologic rest the eye receives from the drug. The surgeon will obtain much assistance and save time by using the retinoscope in cases of aphakia, in old persons especially who are very slow to answer, and will insist upon a description of what they do and do not see, as also in re-reading the test-card from the very top each time a change of lens is put in the trial frame. Presbyopes of fifty or more years of age can be quickly and not inconveniently refracted by the shadow test after having their pupils dilated with a weak (four per cent.) solution of cocain.

Concave Mirror.—While the study of retinoscopy with the concave mirror is not a part of the subject of this book, and allusion to it has been carefully avoided up to this time, yet for the benefit of those who may wish to try it, the writer would suggest that it will be necessary to place the source of light (20 or 30 mm. opening in light-screen) above and beyond the patient's head, one meter distant, or more, so that the convergent rays from the mirror come to a focus and cross before entering the observed eye. Then to estimate the refraction, proceed as with the plane mirror, remembering, however, that the movements of the retinal illumination are just the reverse of those obtained when using the plane mirror.

The Author's Schematic Eye for Studying Retinoscopy.—For illustration see figure 1 and the *Journal of the American Medical Association*, January 5, 1895. The eye as here shown, slightly reduced in size, is made of two brass cylinders, one somewhat smaller than its fellow, to permit slipping evenly into the other. Both cylinders are well blackened inside, and the larger is also blackened outside. The smaller cylinder is closed at one end (concave surface), and on its inner surface is placed a colored lithograph of the normal eye ground. The larger cylinder is also closed at one end, except for a central round opening 10 mm. in diameter, which is occupied by a + 16 D. lens, and on its outer surface is a colored lithograph of the normal eye and its appendages; the pupil is left dilated, and corresponds in size to the central

opening just referred to. In addition to the picture of the eye, there is also lithographed on the upper half of the periphery the degree marks similar to those on a trial frame. To the lower half of the periphery are secured, at equal distances, three posts with grooves to hold trial-lenses. On the side of the small cylinder is an index which records emmetropia, and the amount of myopia and hyperopia, as it is pushed into or drawn out of the large cylinder. The eye is mounted on a convenient stand and upright, so that it may be moved as required. In using this eye, if the red eye ground and retinal vessels disturb the beginner, then he may substitute a piece of white paper for the retina. To study astigmatism with the model, the beginner will have to place a cylinder of known strength in the groove next to the eye and study the characteristic band of light so diagnostic of this condition, and at the same time he should learn to locate the axis of the band with the axonometer.

The author's light-screen or cover chimney (see figure 4 and the *Annals of Ophthalmology and Otology*, October, 1896) is made of one-eighth inch asbestos, and of sufficient size to fit easily over the glass chimney of the Argand burner; attached to the asbestos by means of a metal clamp are two superimposed discs, which revolve independently of each other. The lower disc contains a piece of white porcelain, 30 mm. in diameter; also four round openings, respectively 5, 10, 20, and 35 mm. in diameter. The upper disc contains a round 35 mm. opening, a round section of blue cobalt glass, a perforated disc,

a vertical and a horizontal slit, each 2½ by 25 mm. The several uses of this screen are as follows:

1. For the ophthalmoscope a good light is obtained by superimposing the two 35 mm. openings.

2. Combining the 35 mm. opening in the upper with either the 5 or 10 mm. in the lower disc, a source of light is produced for the small retinoscope; and,

3. By substituting the 20 mm. opening, light is had for the concave mirror.

4. Placing the cobalt glass over the 5, 10, 20, or 35 mm. opening, and the chromo-aberration test for ametropia is given.

5. To test for astigmatism at one meter while using the plane mirror, or for heterophoria at six meters, the perforated disc is to be turned over the porcelain, the latter producing a clear white image.

6. The horizontal slit placed over the porcelain glass, and the operator may exercise the oblique muscles.

7. The vertical slit similarly placed gives the test for paralyzed muscles.

Lenses for the Study of the Scissor Movement, Conic Cornea, Spheric Aberration, and Lenticular Astigmatism.—(Described by the author in the *Journal of the American Medical Association*, December 18, 1897.)

As the scissor movement, conic cornea, spheric aberration, and lenticular astigmatism, as recognized by the retinoscope, are so difficult of demonstration, except in the individual patient, the writer has suggested and had made four lenses which will illustrate

these conditions respectively when placed in front of his schematic eye ; and thus the beginner in retinoscopy may have the opportunity to see, know, and study these important and interesting manifestations (and at small expense) before proceeding direct and in comparative ignorance to his patient.

Figures 45 and 46 represent a plano-concave cylinder of two diopters, mounted in a cell of the trial-case, and to one-half of its plane surface is cemented (at the same axis) a plano-convex cylinder of four diopters, thus making a combination lens, one half

FIG. 45.



FIG. 47.



FIG. 49.

of which is a — 2 D. and the other half equaling a + 2 D. Placing this lens, with its axis at 180°, before the schematic eye at emmetropia (zero), and the observer at one meter distance with his plane mirror, the two light areas characteristic of the scissor movement, with their comparatively straight edges and dark interspace, may be seen approaching each other from above and below (and the dark interspace disappearing) as the mirror is tilted in the vertical meridian.

Figures 47 and 48 represent a section of thin plane glass mounted as in figure 45, and has cemented at

its center a small plano-convex sphere of three diopters, whose base is about four mm. in diameter. Placing this lens in front of the schematic eye at emmetropia, and reflecting the light from the plane mirror at one meter, there will be seen in the pupillary area a small central illumination, which moves against or opposite to the movement of the mirror, and at the same time there will also be seen a peripheral ring (at the edge of the iris) which moves rapidly with the movement of the mirror; between these light areas is a shaded ring of feeble illumination. This



FIG. 46.

FIG. 48.

is the retinoscopic picture and movement of the light areas, so indicative of conic cornea. It is also an exaggerated picture of negative aberration.

Figures 40 and 50 represent a section similar to that shown in figures 47 and 48, except that at its center is ground a — 2 D. sphere of about four mm. in diameter. To produce spheric aberration of the positive form, place this lens in front of the schematic eye at emmetropia, and the observer, seated at one meter distance with the plane mirror,

will see in the pupillary area a central illumination which moves slower than the peripheral area or ring (at the edge of the iris), which moves rapidly, both areas moving with the movement of the mirror. Figure 51 shows a lens for studying lenticular astigmatism. This is made by scratching a piece of plane glass with a diamond.

After the observer has carefully studied these pictures, it will be obvious that changes other than those mentioned can be made with these lenses, and he should proceed to note them by—

FIG. 50.

FIG. 51.

1. Changing the focus of the schematic eye.

2. By varying his distance from the eye.

3. By placing both the concave and convex spheres in combination.

4. By placing a concave cylinder in front of the double cylinder at an oblique axis, thus getting a picture of compound irregular astigmatism.

5. By placing a concave cylinder in front of the convex sphere and developing astigmatism with the conic cornea, which is the usual condition; or a con-

vex cylinder might be used in place of the concave cylinder if a higher error is desired.

6. It is obvious, also, that the scissor movement can be produced by a prism which is made to cover one-half of the pupillary area, but the resulting picture is not so satisfactory for demonstration as that given by the combination lens referred to in figure 45.

INDEX.

ABERRATION, negative, 77, 78, 79
positive, 77, 78, 79
spheric, 77, 77, 78
Accommodation, 29
cramp of, 75
Accuracy, 23, 41
advantages of, 20, 21
Albino, 37
Amblyopia, 21
Aphakia, 21, 75
Apparatus, 21, 47, 48, 49, 50, 51
Area of light, 38, 77, 78
Argand burner, 26
Arrangement of light, 32, 33
of observer, 32, 33
of patient, 32, 33
Astigmatism, 46, 48, 60, 62, 63
compound irregular, 75, 76
corneal, 70
irregular, 70
lenticular, 70, 71
regular, 60, 61, 64, 65
Author's axonometer, 30
retinoscope, 25, 35
schematic eye, 22, 23, 80, 81
schematic lenses, 82, 83, 84, 85, 86
shade, 26, 27, 81, 82
trial-case, 51
trial-frame, 50
Avoid, what to, 39
Axiom, 21
Axonometer, celluloid, 30, 50, 69
metal, 30, 66, 67

BAND of light, 46, 61, 62, 63
Bands of light, 72, 73, 74
causes of, 74, 75
Beginner, 23, 36
Brunette, 37
Burner, Argand, 26

CATOPTRIC images, 37, 44
Central shadow, 24, 25, 29, 44
illumination, 38
Children, 21
Cocain, 79
Compound irregular astigmatism, 75, 76
Concave mirror, 19, 28, 80, 82
Conic cornea, 76, 77, 78
astigmatism in, 70
Conjugate focus, 23, 40
Cornea, apex of, 39, 51
Cover chimney, 26, 27, 81, 82
Crossed cylinders, 66
Cycloplegic, 29, 39, 71, 79

DARKNESS, 38, 39, 44
Dark room, 27, 28
Definition, 19
Dioptroscopy, 19
Direction of movement, 39, 40, 41, 42, 45
Discs, 24, 25, 35, 48, 49
Distance, 31

EMMETROPIA, 41, 45, 55, 56
Examples, 41, 53

FACIAL illumination, 41, 52
Fantoscopy, 19
Far-point, 40, 58
Feeble-minded, 21
Flame, 26, 27
Form of illumination, 39, 40, 46
Formula, 61
Fundus-reflex test, 19, 62

GENERAL appearances, 36, 37

HOW to use the mirror, 34, 35
Hyaloid vessel, 43
Hyperopia, 41, 45, 46, 52, 53, 54
Hyperopic astigmatism, 60, 61

ILLITERATES, 21
Illuminated area, 38
Illumination, facial, 41, 52
form of, 38
retinal, 38, 39, 43, 52
Illustrations, 22, 25, 26, 27, 32, 33, 34, 35, 38, 46, 47, 48, 50, 51, 52, 53, 54, 55, 56, 57, 58, 61, 62, 63, 64, 67, 69, 71, 72, 73, 74, 75, 77, 78, 83, 84, 85
Image, 38
Images, 37
Inaccuracy, 21
Irregular astigmatism, 70, 71
corneal, 77
lenticular, 70, 71

JACKSON, 24
Jennings, 48, 49

KALEIDOSCOPE, 70
Keratoscopy, 19
Koroscopy, 19

LAMP, electric, 26
gas, 26
oil, 26
Lenses, neutralizing, 37
rule for, 40
schematic, 46, 47
Lenticular astigmatism, 70, 71
Light, 25
electric, 16
gas, 26
oil, 26
Welsbach, 25, 26
Light-screen, 26, 27

MACULA, 30, 56
Macular region, 30
Magnification, point of, 40
Meter distance, 33
stick, 31
Mirror, 24, 25, 35, 80
Mixed astigmatism, 64, 65
Movement of light, 39
in pupillary area, 44, 45, 52
on face, 39
of mirror, 35, 39, 47
Mulatto, 37
Myopia, 45, 46, 56, 57, 58

NAME, 19
Negative aberration, 77, 78, 79
Neutralizing lenses, 43, 44
rules for, 46, 47, 58, 59
Nystagmus, 21

OBSERVER, 29
Oil-lamp, 26
Ophthalmoscope, 20, 60
Optician, 20

PATIENT, 29, 30
Pearse, 66
Point of magnification, 40
of reversal, 20, 37, 40, 42, 43, 54, 56, 57, 58
to find, 40, 43
Position of light, 28
of mirror, 28
of observer, 29
of patient, 29
of lenses, 44
Positive aberration, 77, 78, 79
Post-cycloplegic, 71
Principle of retinoscopy, 20, 40
Punctum remotum, 40, 58
Pupillary area, 36, 42, 70
Pupilloscopy, 19

QUICK movement, 47, 48

RATE of movement of illumination, 39, 45, 46
of mirror, 45, 46
Reflection from cornea, 36, 37, 34
from lenses, 37, 44
from mirror, 34
Refraction at center of pupil, 39, 51
at pupillary edge, 37
Regular astigmatism, 60, 61, 64, 65

Retinal illumination, 38, 39, 43
image, 39
vessel, 43, 44
Retinophotoscopy, 19
Retinoscope, 24, 25, 35
Retinoscopy, 14, 19, 20
advantages of, 21
in amblyopia, 21
in children, 21
in feeble-minded, 21
in nystagmus, 21
without a cycloplegic, 79
Retinoskiascopy, 19
Reversal of movement, 20, 37, 40, 42, 43, 45, 46, 54, 56, 57, 58
Room, 27, 28
Rule, Pearse's, 66
Rules for distance, 58, 59
for lenses, 46, 47, 58, 59

SCHEMATIC eye, 22
lenses, 46, 47
Scissor movement, 72, 73, 74
Shade, 26, 27
Shadow, 38, 39, 44
test, 38
Sight-hole, 24, 44
Size of mirror, 24, 25
of sight-hole, 24, 25
Skiagraphy, 14
Skiascopy, 14
Slow movement, 47
Source of light, 32, 33
Spheric aberration, 37
Squint, 30
use of axonometer in cases of, 30
Suggestions to the beginner, 23

THORINGTON, 22, 23, 25, 26, 27, 30, 35, 50, 51
Trial-case, 51
-frame, 50, 51

UMBRASCOPY, 19

VALUE of retinoscopy, 20
Vision of observer, 29
of patient, 30

WELSBACH, 25, 26
What the observer sees, 36, 37
to avoid, 39, 40, 43, 44
Where to look, and what to look for, 39
Würdemann, 47, 49

YOUNG children, 21

No. 8 November, 1900

A Classified Catalogue of Books on Medicine and the Collateral Sciences, Pharmacy, Dentistry, Chemistry, Hygiene, Microscopy, Etc.

P. Blakiston's Son & Company, Publishers of Medical and Scientific Books,
1012 Walnut Street, Philadelphia

SUBJECT INDEX.

Special Catalogues of Books on Pharmacy, Dentistry, Chemistry, Hygiene, and Nursing will be sent free upon application. All inquiries regarding prices, dates of edition, terms, etc., will receive prompt attention.

SUBJECT	PAGE
Alimentary Canal (see Surgery)	19
Anatomy	3
Anesthetics	14
Autopsies (see Pathology)	16
Bacteriology (see Pathology)	16
Bandaging (see Surgery)	19
Blood, Examination of	16
Brain	4
Chemistry	4
Children, Diseases of	6
Climatology	14
Clinical Charts	6
Compends	22, 23
Consumption (see Lungs)	11
Cyclopedia of Medicine	8
Dentistry	7
Diabetes (see Urin. Organs)	21
Diagnosis	17
Diagrams (see Anatomy)	3
Dictionaries, Cyclopedias	8
Diet and Food	14
Dissectors	3
Ear	9
Electricity	9
Emergencies (see Surgery)	19
Eye	9
Fevers	9
Gout	10
Gynecology	21
Hay Fever	20
Heart	10
Histology	10
Hospitals (see Hygiene)	11
Hydrotherapy	14
Hygiene	11
Insanity	4
Intestines (see Miscellaneous)	14
Latin, Medical (see Miscellaneous and Pharmacy)	14, 16
Life Insurance	14
Lungs	12
Massage	12
Materia Medica	12
Medical Jurisprudence	13
Microscopy	13
Milk Analysis (see Chemistry)	4
Miscellaneous	14
Nervous Diseases	14
Nose	20
Nursing	15
Obstetrics	16
Ophthalmology	9
Organotherapy	14
Osteology (see Anatomy)	3
Pathology	16
Pharmacy	16
Physical Diagnosis	17
Physical Training (see Miscellaneous)	14
Physiology	17
Pneumotherapy	14
Poisons (see Toxicology)	13
Popular Medicine	10
Practice of Medicine	18
Prescription Books	18
Refraction (see Eye)	9
Rheumatism	10
Sanitary Science	11
Skin	19
Spectacles (see Eye)	9
Spine (see Nervous Diseases)	14
Stomach (see Miscellaneous)	14
Students' Compends	22, 23
Surgery and Surgical Diseases	19
Syphilis	21
Technological Books	4
Temperature Charts	6
Therapeutics	12
Throat	20
Toxicology	13
Tumors (see Surgery)	19
U. S. Pharmacopœia	17
Urinary Organs	20
Urine	20
Venereal Diseases	21
Veterinary Medicine	21
Visiting Lists, Physicians'. (*Send for Special Circular.*)	
Water Analysis	11
Women, Diseases of	21

☞ *The prices as given in this Catalogue are net. Cloth binding, unless otherwise specified. Postpaid, upon receipt of advertised price.*

SPECIAL NOTE.—The prices given in this catalogue are absolutely net, no discount can be allowed retail purchasers under any consideration. This rule has been established in order that everyone will be treated alike, a general reduction in former prices having been made to meet previous retail discounts. Upon receipt of the advertised price any book will be forwarded by mail or express, **all charges prepaid.**

ANATOMY.

MORRIS. Text-Book of Anatomy. 2d Edition. Revised and Enlarged. 790 Illustrations, 214 of which are printed in colors. Cloth, $6.00; Leather, $7.00; Half Russia, $8.00

"The ever-growing popularity of the book with teachers and students is an index of its value, and it may safely be recommended to all interested."—*Medical Record, New York.*

☞ Sample pages and illustrations will be sent free to any address.

BROOMELL. Anatomy and Histology of the Human Mouth and Teeth. 284 Illustrations. $4 50

DEAVER. Surgical Anatomy. A Treatise on Human Anatomy in its Application to Medicine and Surgery. With about 400 very Handsome full-page Illustrations Engraved from Original Drawings made by special Artists from dissections prepared for the purpose. Three Volumes. Royal Square Octavo.
Cloth, $21 00; Half Morocco or Sheep, $24.00; Half Russia, $27 00

ECKLEY. Practical Anatomy. A Manual for use in the Dissecting Room. Based upon Morris' Text-Book of Anatomy and Including a Section on the Fundamental Principles of Anatomy. With 347 Illustrations, many of which are in colors. Cloth, $3 50; Oil Cloth, $4.00

GORDINIER. Anatomy of the Central Nervous System. With 271 Illustrations, many of which are original. Cloth, $6.00

HEATH. Practical Anatomy. 8th Edition. 300 Illus. $4.25

HOLDEN. Anatomy. A Manual of the Dissections of the Human Body. Carefully Revised by A. HEWSON, M.D., Demonstrator of Anatomy, Jefferson Medical College. Philadelphia. Over 300 handsome Illustrations. 7th Edition. In two compact 12mo Volumes, Large New Type. Bound in Oil Cloth, price of each Volume, $1.50

HOLDEN. Human Osteology. Comprising a Description of the Bones, with Colored Delineations of the Attachments of the Muscles. The General and Microscopical Structure of Bone and its Development. With Lithographic Plates and numerous Illus. 8th Ed. $5.25

HOLDEN. Landmarks. Medical and Surgical. 4th Ed. .75

MACALISTER. Human Anatomy. Systematic and Topographical, including the Embryology, Histology, and Morphology of Man. With Special Reference to the Requirements of Practical Surgery and Medicine. 816 Illustrations. Cloth, $5.00; Leather, $6.00

MARSHALL. Physiological Diagrams. Life Size. Colored. Eleven Life-Size Diagrams (each seven feet by three feet seven inches). Designed for Demonstration before the Class.
In Sheets, Unmounted, $40.00; Backed with Muslin and Mounted on Rollers, $60.00; Ditto, Spring Rollers, in Handsome Walnut Wall Map Case, $100.00; Single Plates—Sheets, $5.00; Mounted, $7.50. Explanatory Key, .50. *Purchaser must pay freight charges.*

POTTER. Compend of Anatomy, Including Visceral Anatomy. 6th Ed. 16 Lith. Plates and 117 other Illus. .80; Interleaved, $1.00

WILSON. Anatomy. 11th Edition. 429 Illus., 26 Plates. $5.00

WINDLE. Surface Anatomy. Colored and other Illus. $1.00

BRAIN AND INSANITY (see also Nervous Diseases).

BLACKBURN. A Manual of Autopsies. Designed for the Use of Hospitals for the Insane and other Public Institutions. Ten full-page Plates and other Illustrations. $1.25

GORDINIER. The Gross and Minute Anatomy of the Central Nervous System. With many full-page and other Illustrations. 8vo. Cloth, $6.00; Sheep, $7.00

HORSLEY. The Brain and Spinal Cord. The Structure and Functions of. Numerous Illustrations. $2.50

LEWIS (BEVAN). Mental Diseases. A Text-Book Having Special Reference to the Pathological Aspects of Insanity. 26 Lithographic Plates and other Illustrations. 2d Ed. *Just Ready.* $7.00

MANN. Manual of Psychological Medicine and Allied Nervous Diseases. $3.00

REGIS. Mental Medicine. Authorized Translation by H. M. BANNISTER, M.D. $2.00

SHUTTLEWORTH. Mentally Deficient Children. New Edition. $1.50

STEARNS. Mental Diseases. With a Digest of Laws Relating to Care of Insane. Illustrated. Cloth, $2.75; Sheep, $3.25

TUKE. Dictionary of Psychological Medicine. Giving the Definition, Etymology, and Symptoms of the Terms used in Medical Psychology, with the Symptoms, Pathology, and Treatment of the Recognized Forms of Mental Disorders. Two volumes. $10.00

WOOD, H. C. Brain and Overwork. .40

CHEMISTRY AND TECHNOLOGY.

Special Catalogue of Chemical Books sent free upon application.

ALLEN. Commercial Organic Analysis. A Treatise on the Modes of Assaying the Various Organic Chemicals and Products Employed in the Arts, Manufactures, Medicine, etc., with concise methods for the Detection of Impurities, Adulterations, etc. 8vo.

Vol. I. Alcohols, Neutral Alcoholic Derivatives, etc., Ethers, Vegetable Acids, Starch, Sugars, etc. 3d Edition, by HENRY LEFFMANN, M. D. $4.50

Vol. II, Part I. Fixed Oils and Fats, Glycerol, Explosives, etc. 3d Edition, by HENRY LEFFMANN, M. D. $3.50

Vol. II, Part II. Hydrocarbons, Mineral Oils, Lubricants, Benzenes, Naphthalenes and Derivatives, Creosote, Phenols, etc. 3d Edition, by HENRY LEFFMANN, M.D. $3.50

Vol. II, Part III. Terpenes, Essential Oils, Resins, Camphors, etc. 3d Edition, by HENRY LEFFMANN, M.D. *Preparing.*

Vol. III, Part I. Tannins, Dyes and Coloring Matters. 3d Edition. Enlarged and Rewritten by J. MERRITT MATTHEWS, PH.D., of the Philadelphia Textile School. Illustrated. $4.50

Vol. III, Part II. The Amines, Hydrazines and Derivatives, Pyridine Bases. The Antipyretics, etc. Vegetable Alkaloids, Tea, Coffee, Cocoa, etc. 8vo. 2d Edition. $4.50

Vol. III, Part III. Vegetable Alkaloids, Non-Basic Vegetable Bitter Principles. Animal Bases, Animal Acids, Cyanogen Compounds, etc. 2d Edition, 8vo. $4.50

Vol. IV. The Proteids and Albuminous Principles. 2d Ed. $4.50

ALLEN. Albuminous and Diabetic Urine. Illustrated. $2.25

BARTLEY. Medical and Pharmaceutical Chemistry. A Text-Book for Medical, Dental, and Pharmaceutical Students. With Illustrations, Glossary, and Complete Index. 5th Edition, carefully Revised. Cloth, $3.00; Sheep, $3.50

BARTLEY. Clinical Chemistry. The Examination of Feces, Saliva, Gastric Juice, Milk, and Urine. $1.00

BLOXAM. Chemistry, Inorganic and Organic. With Experiments. 8th Ed., Revised. 281 Engravings. Clo., $4.25; Lea., $5.25

CALDWELL. Elements of Qualitative and Quantitative Chemical Analysis. 3d Edition, Revised. $1.00

CAMERON. Oils and Varnishes. With Illustrations. $2.25

CAMERON. Soap and Candles. 54 Illustrations $2.00

CLOWES AND COLEMAN. Quantitative Analysis. 5th Edition. 122 Illustrations. *Just Ready.* $3.50

CONGDON. Laboratory Instructions in Chemistry. *In Press.*

GARDNER. The Brewer, Distiller, and Wine Manufacturer. Illustrated. $1.50

GARDNER. Bleaching, Dyeing, and Calico Printing. $1.50

GROVES AND THORP. Chemical Technology. The Application of Chemistry to the Arts and Manufactures.

Vol. I. Fuel and Its Applications. 607 Illustrations and 4 Plates. Cloth, $5.00; ½ Mor, $6.50
Vol. II. Lighting. Illustrated. Cloth, $4.00; ½ Mor., $5.50
Vol. III. Gas Lighting. Cloth, $3.50; ½ Mor., $4.50
Vol. IV. Electric Lighting. *In Press.*

HOLLAND. The Urine, the Gastric Contents, the Common Poisons, and the Milk. Memoranda, Chemical and Microscopical, for Laboratory Use. 6th Ed. Illustrated and interleaved, $1.00

LEFFMANN. Compend of Medical Chemistry, Inorganic and Organic. Including Urine Analysis. 4th Edition, Rewritten and Revised. .80; Interleaved, $1.00

LEFFMANN. Analysis of Milk and Milk Products. 2d Edition, Enlarged. Illustrated. $1.25

LEFFMANN. Water Analysis. For Sanitary and Technic Purposes. Illustrated. 4th Edition. $1.25

LEFFMANN. Structural Formulæ. Including 180 Structural and Stereo-Chemical Formulæ. 12mo. Interleaved. $1.00

MUTER. Practical and Analytical Chemistry. 2d American from the Eighth English Edition Revised to meet the requirements of American Students. 56 Illustrations. $1.25

OETTEL. Exercises in Electro-Chemistry. Illustrated. .75

OETTEL. Electro-Chemical Experiments. Illustrated. .75

RICHTER. Inorganic Chemistry. 5th American from 10th German Edition. Authorized translation by EDGAR F. SMITH, M.A., PH.D. 89 Illustrations and a Colored Plate. $1.75

RICHTER. Organic Chemistry. 3d American Edition. Trans. from the 8th German by EDGAR F. SMITH. Illustrated. 2 Volumes.

Vol. I. Aliphatic Series. 625 Pages. $3.00
Vol. II. Carbocyclic Series. 671 Pages. $3.00

SMITH. Electro-Chemical Analysis. 2d Edition, Revised. 28 Illustrations. $1.25

SMITH AND KELLER. Experiments. Arranged for Students in General Chemistry. 4th Edition. Illustrated. *Just Ready.* .60

SUTTON. Volumetric Analysis. A Systematic Handbook for the Quantitative Estimation of Chemical Substances by Measure, Applied to Liquids, Solids, and Gases. 8th Edition, Revised. 112 Illustrations. *Just Ready.* $5.00

SYMONDS. Manual of Chemistry, for Medical Students. 2d Edition. $2.00

TRAUBE. Physico-Chemical Methods. Translated by Hardin. 97 Illustrations. $1.50

THRESH. Water and Water Supplies. 2d Edition. $2.00

ULZER AND FRAENKEL. Chemical Technical Analysis. Translated by Fleck. Illustrated. $1.25

WOODY. Essentials of Chemistry and Urinalysis. 4th Edition. Illustrated. $1.50

*** *Special Catalogue of Books on Chemistry free upon application.*

CHILDREN.

CAUTLEY. Feeding of Infants and Young Children by Natural and Artificial Methods. $2.00

HALE. On the Management of Children. .50

HATFIELD. Compend of Diseases of Children. With a Colored Plate. 2d Edition. .80; Interleaved, $1.00

MEIGS. Infant Feeding and Milk Analysis. The Examination of Human and Cow's Milk, Cream, Condensed Milk, etc., and Directions as to the Diet of Young Infants. .50

POWER. Surgical Diseases of Children and their Treatment by Modern Methods. Illustrated. $2.50

SHUTTLEWORTH. Mentally Deficient Children. New Edition. $1.50

STARR. The Digestive Organs in Childhood. The Diseases of the Digestive Organs in Infancy and Childhood. With Chapters on the Investigation of Disease and the Management of Children. 3d Edition, Enlarged. Illustrated by two Colored Plates and numerous Wood Engravings. *In Preparation.*

STARR. Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

SMITH. Wasting Diseases of Children. 6th Edition. $2.00

TAYLOR AND WELLS. The Diseases of Children. Illustrated. A New Manual. 746 pages. $4 00

CLINICAL CHARTS.

GRIFFITH. Graphic Clinical Chart for Recording Temperature, Respiration, Pulse, Day of Disease, Date, Age, Sex, Occupation, Name, etc. Printed in three colors. Sample copies free. Put up in loose packages of fifty, .50. Price to Hospitals, 500 copies, $4.00; 1000 copies, $7.50. With name of Hospital printed on, .50 extra.

KEEN'S CLINICAL CHARTS. Seven Outline Drawings of the Body, on which may be marked the Course of Disease, Fractures, Operations, etc. Pads of fifty, $1.co. Each Drawing may also be had separately, twenty-five to pad, 25 cents.

SCHREINER. Diet Lists. Arranged in the form of a chart. With Pamphlets of Specimen Dietaries. Pads of 50. .75

DENTISTRY.

Special Catalogue of Dental Books sent free upon application.

BARRETT. Dental Surgery for General Practitioners and Students of Medicine and Dentistry. Extraction of Teeth, etc. 3d Edition. Illustrated. *Nearly Ready.*

BROOMELL. Anatomy and Histology of the Human Mouth and Teeth. 284 Handsome Illustrations. $4.50

FILLEBROWN. A Text-Book of Operative Dentistry. Written by invitation of the National Association of Dental Faculties. Illustrated. $2.25

GORGAS. Dental Medicine. A Manual of Materia Medica and Therapeutics. 6th Edition, Revised. Cloth, $4.00; Sheep, $5.00

HARRIS. Principles and Practice of Dentistry. Including Anatomy, Physiology, Pathology, Therapeutics, Dental Surgery, and Mechanism. 13th Edition. Revised by F. J. S. GORGAS, M.D., D.D.S. 1250 Illustrations. Cloth, $6.00; Leather, $7.00

HARRIS. Dictionary of Dentistry. Including Definitions of Such Words and Phrases of the Collateral Sciences as Pertain to the Art and Practice of Dentistry. 6th Edition. Revised and Enlarged by FERDINAND F. S. GORGAS, M.D., D.D.S. Cloth, $5.00; Leather, $6.00

HEATH. Injuries and Diseases of the Jaws. 4th Edition. 187 Illustrations. $4.50

HEATH. Lectures on Certain Diseases of the Jaws. 64 Illustrations. Boards, .50

RICHARDSON. Mechanical Dentistry. 7th Edition. Thoroughly Revised and Enlarged by DR. GEO. W. WARREN. 691 Illustrations. Cloth, $5.00; Leather, $6.00

SMITH. Dental Metallurgy. Illustrated. $1.75

TAFT. Index of Dental Periodical Literature. $2.00

TALBOT. Irregularities of the Teeth and Their Treatment. 2d Edition. 234 Illustrations. $3.00

TOMES. Dental Anatomy. Human and Comparative. 263 Illustrations. 5th Edition. $4.00

TOMES. Dental Surgery. 4th Edition. 289 Illustrations. $4.00

WARREN. Compend of Dental Pathology and Dental Medicine. With a Chapter on Emergencies. 3d Edition. Illustrated. .80; Interleaved, $1.25

WARREN. Dental Prosthesis and Metallurgy. 129 Ills. $1.25

WHITE. The Mouth and Teeth. Illustrated. .40

*** *Special Catalogue of Dental Books free upon application.*

DICTIONARIES.

GOULD. The Illustrated Dictionary of Medicine, Biology, and Allied Sciences. Being an Exhaustive Lexicon of Medicine and those Sciences Collateral to it: Biology (Zoology and Botany), Chemistry, Dentistry, Parmacology, Microscopy, etc., with many useful Tables and numerous fine Illustrations. 1633 pages. 5th Ed.
Sheep or Half Dark Green Leather, $10.00; Thumb Index, $11.00
Half Russia, Thumb Index, $12.00

GOULD. The Medical Student's Dictionary. 11th Edition. Illustrated. Including all the Words and Phrases Generally Used in Medicine, with their Proper Pronunciation and Definition, Based on Recent Medical Literature. With a new Table of Eponymic Terms and Tests and Tables of the Bacilli, Micrococci, Mineral Springs, etc., of the Arteries, Muscles, Nerves, Ganglia, Plexuses, etc. **11th Edition.** Enlarged by over 100 pages and illustrated with a large number of engravings. 840 pages. *Just Ready.*
Half Green Morocco, $2.50; Thumb Index, $3 00

GOULD. The Pocket Pronouncing Medical Lexicon. 4th Edition. (30,000 Medical Words Pronounced and Defined.) Containing all the Words, their Definition and Pronunciation, that the Medical, Dental, or Pharmaceutical Student Generally Comes in Contact With; also Elaborate Tables of Eponymic Terms, Arteries, Muscles, Nerves, Bacilli, etc., etc., a Dose List in both English and Metric Systems, etc., Arranged in a Most Convenient Form for Reference and Memorizing. **A new (Fourth) Edition, Revised and Enlarged.** 838 pages. *Just Ready.*
Full Limp Leather, Gilt Edges, $1.00; Thumb Index, $1.25

100,000 Copies of Gould's Dictionaries Have Been Sold.

GOULD AND PYLE. Cyclopedia of Practical Medicine and Surgery. Seventy-two Special Contributors. Illustrated. One Volume. A Concise Reference Handbook, Alphabetically Arranged, of Medicine, Surgery, Obstetrics, Materia Medica, Therapeutics, and the Various Specialties, with Particular Reference to Diagnosis and Treatment. Compiled under the Editorial Supervision of GEORGE M. GOULD, M.D., Author of "An Illustrated Dictionary of Medicine"; Editor "Philadelphia Medical Journal," etc.; and WALTER L. PYLE, M.D., Assistant Surgeon Wills Eye Hospital; formerly Editor "International Medical Magazine," etc., and Seventy-two Special Contributors. With many Illustrations. Large Square 8vo, to correspond with Gould's "Illustrated Dictionary." *Just Ready.* Full Sheep or Half Dark-Green Leather, $10.00
With Thumb Index, $11.00; Half Russia, Thumb Index, $12.00 net.

*** Sample Pages and Illustrations and Descriptive Circulars of Gould's Dictionaries and Cyclopedia sent free upon application.

HARRIS. Dictionary of Dentistry. Including Definitions of Such Words and Phrases of the Collateral Sciences as Pertain to the Art and Practice of Dentistry. 6th Edition. Revised and Enlarged by FERDINAND J. S. GORGAS, M.D., D.D.S. Cloth, $5.00; Leather, $6 00

LONGLEY. Pocket Medical Dictionary. With an Appendix, containing Poisons and their Antidotes, Abbreviations used in Prescriptions, etc. Cloth, .75; Tucks and Pocket, $1.00

MAXWELL. Terminologia Medica Polyglotta. By Dr. THEODORE MAXWELL, Assisted by Others. $3.00
The object of this work is to assist the medical men of any nationality in reading medical literature written in a language not their own. Each term is usually given in seven languages, viz.: English, French, German, Italian, Spanish, Russian, and Latin.

TREVES AND LANG. German-English Medical Dictionary.
Half Russia, $3.25

EAR (see also Throat and Nose).

BURNETT. Hearing and How to Keep It. Illustrated. .40

DALBY. Diseases and Injuries of the Ear. 4th Edition. 38 Wood Engravings and 8 Colored Plates. $2.50

HOVELL. Diseases of the Ear and Naso-Pharynx. Including Anatomy and Physiology of the Organ, together with the Treatment of the Affections of the Nose and Pharynx which Conduce to Aural Disease. 122 Illustrations. 2d Edition. *Preparing.*

PRITCHARD. Diseases of the Ear. 3d Edition, Enlarged. Many Illustrations and Formulæ. $1.50

WOAKES. Deafness, Giddiness, and Noises in the Head. 4th Edition. Illustrated. $2.00

ELECTRICITY.

BIGELOW. Plain Talks on Medical Electricity and Batteries. With a Therapeutic Index and a Glossary. 43 Illustrations. 2d Edition. $1.00

HEDLEY. Therapeutic Electricity and Practical Muscle Testing. 99 Illustrations. *Just Ready.* $2.50

JACOBI. Electrotherapy. Illustrated. *In Press.*

JONES. Medical Electricity. 3d Edition. 117 Illus. $3 00

MASON. Medical Electricity. Numerous Illustrations. .75

EYE.

A Special Circular of Books on the Eye sent free upon application

DONDERS. The Nature and Consequences of Anomalies of Refraction. With Portrait and Illustrations. Half Morocco, $1.25

FICK. Diseases of the Eye and Ophthalmoscopy. Translated by A. B. HALE, M. D. 157 Illustrations, many of which are in colors, and a glossary. Cloth, $4.50; Sheep, $5.50

GOULD AND PYLE. Compend of Diseases of the Eye and Refraction. Including Treatment and Operations, and a Section on Local Therapeutics. With Formulæ, Useful Tables, a Glossary, and 111 Illus., several of which are in colors. 2d Edition, Revised. Cloth, .80; Interleaved, $1.00

GOWERS. Medical Ophthalmoscopy. A Manual and Atlas with Colored Autotype and Lithographic Plates and Wood-cuts, Comprising Original Illustrations of the Changes of the Eye in Diseases of the Brain, Kidney, etc. 3d Edition. $4.00

HARLAN. Eyesight, and How to Care for It. Illus. .40

HARTRIDGE. Refraction. 104 Illustrations and Test Types. 10th Edition, Enlarged. *Just Ready.* $1.50

HARTRIDGE. On the Ophthalmoscope. 3d Edition. With 4 Colored Plates and 68 Wood-cuts. $1.50

HANSELL AND REBER. Muscular Anomalies of the Eye. Illustrated. $1.50

HANSELL AND BELL. Clinical Ophthalmology. Colored Plate of Normal Fundus and 120 Illustrations. $1.50

JESSOP. Manual of Ophthalmic Surgery and Medicine. Colored Plates and 108 other Illustrations. Cloth, $3.00

MORTON. Refraction of the Eye. Its Diagnosis and the Correction of its Errors. 6th Edition. $1.00

OHLEMANN. Ocular Therapeutics. Authorized Translation, and Edited by DR. CHARLES A. OLIVER. $1 75

PHILLIPS. Spectacles and Eyeglasses. Their Prescription and Adjustment. 2d Edition. 49 Illustrations. $1.00

SWANZY. Diseases of the Eye and Their Treatment. 7th Edition, Revised and Enlarged. 164 Illustrations, 1 Plain Plate, and a Zephyr Test Card. *Just Ready.* $2.50

THORINGTON. Retinoscopy. 3d Edition. Illustrated. $1.00

THORINGTON. Refraction and How to Refract. 200 Illustrations, 13 of which are Colored. 2d Edition. *Just Ready.* $1.50

WALKER. Students' Aid in Ophthalmology. Colored Plate and 40 other Illustrations and Glossary. $1.50

WRIGHT. Ophthalmology. 2d Edition, Revised and Enlarged. With new Illustrations. *In Press.*

FEVERS.

COLLIE. On Fevers. Their History, Etiology, Diagnosis, Prognosis, and Treatment. Colored Plates. $2.00

GOODALL AND WASHBOURN. Fevers and Their Treatment. Illustrated. $3.00

GOUT AND RHEUMATISM.

DUCKWORTH. A Treatise on Gout. With Chromo-lithographs and Engravings. Cloth, $6.00

HAIG. Causation of Disease by Uric Acid. A Contribution to the Pathology of High Arterial Tension, Headache, Epilepsy, Gout, Rheumatism, Diabetes, etc. 5th Edition. *Just Ready.* $3 00

HEART.

SANSOM. Diseases of the Heart. The Diagnosis and Pathology of Diseases of the Heart and Thoracic Aorta. With Plates and other Illustrations. $6.00

THORNE. The Schott Methods of the Treatment of Chronic Heart Disease. Third Edition. Illustrated. *Just Ready.* $1 75

HISTOLOGY.

STIRLING. Outlines of Practical Histology. 368 Illustrations. 2d Edition, Revised and Enlarged. With new Illustrations. $2.00

STÖHR. Histology and Microscopical Anatomy. Translated and Edited by A. SCHAPER, M.D., Harvard Medical School. Second American from 7th German Edition, Revised and Enlarged. 292 Illustrations. $3.00

HYGIENE AND WATER ANALYSIS.

Special Catalogue of Books on Hygiene sent free upon application.

CANFIELD. Hygiene of the Sick-Room. A Book for Nurses and Others. Being a Brief Consideration of Asepsis, Antisepsis, Disinfection, Bacteriology, Immunity, Heating, Ventilation, etc. $1.25

COPLIN. Practical Hygiene. A Complete American Text-Book. 138 Illustrations. New Edition. *Preparing.*

ERNST. Prophylaxis and Personal Hygiene. *In Press.*

HARTSHORNE. Our Homes. Illustrated. .40

KENWOOD. Public Health Laboratory Work. 116 Illustrations and 3 Plates. $2.00

LEFFMANN. Examination of Water for Sanitary and Technical Purposes. 4th Edition. Illustrated. $1.25

LEFFMANN. Analysis of Milk and Milk Products. Illustrated. Second Edition $1.25

LINCOLN. School and Industrial Hygiene. .40

McNEILL. The Prevention of Epidemics and the Construction and Management of Isolation Hospitals. Numerous Plans and Illustrations. $3.50

NOTTER. The Theory and Practice of Hygiene. 15 Plates and 138 other Illustrations. 8vo. 2d Edition. *Just Ready.* $7.00

PARKES. Hygiene and Public Health. By Louis C. Parkes, M.D. 5th Edition. Enlarged. Illustrated. $2.50

PARKES. Popular Hygiene. The Elements of Health. A Book for Lay Readers. Illustrated. $1.25

STARR. The Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

STEVENSON AND MURPHY. A Treatise on Hygiene. By Various Authors. In Three Octave Volumes. Illustrated. Vol. I, $6.00; Vol. II, $6.00; Vol. III, $5.00

*** Each Volume sold separately. Special Circular upon application.

THRESH. Water and Water Supplies. 2d Edition. $2.00

WILSON. Hand-Book of Hygiene and Sanitary Science. With Illustrations. 8th Edition. $3.00

WEYL. Sanitary Relations of the Coal-Tar Colors. Authorized Translation by HENRY LEFFMANN, M.D., PH.D. $1.25

LUNGS AND PLEURÆ.

HARRIS AND BEALE. Treatment of Pulmonary Consumption. $2.50

KNOPF. Pulmonary Tuberculosis. Its Modern Prophylaxis and Treatment in Special Institutions and at Home. Illus. $3.00

POWELL. Diseases of the Lungs and Pleuræ, including Consumption. Colored Plates and other Illus. 4th Ed. $4.00

MASSAGE.

KLEEN. Hand-Book of Massage. Authorized translation by MUSSEY HARTWELL, M.D., PH.D. With an Introduction by Dr. S. WEIR MITCHELL. Illustrated by a series of Photographs Made Especially by DR. KLEEN for the American Edition. $2.25

OSTROM. Massage and the Original Swedish Movements. Their Application to Various Diseases of the Body. A Manual for Students, Nurses, and Physicians. Fourth Edition, Enlarged. 105 Illustrations, many of which are original. $1.00

MITCHELL AND GULICK. Mechanotherapy. Illus. *In Press.*

WARD. Notes on Massage. Interleaved. Paper cover, $1.00

MATERIA MEDICA AND THERAPEUTICS.

BIDDLE. Materia Medica and Therapeutics. Including Dose List, Dietary for the Sick, Table of Parasites, and Memoranda of New Remedies. 13th Edition, Revised. 64 Illustrations and a Clinical Index. Cloth, $4.00; Sheep, $5.00

BRACKEN. Outlines of Materia Medica and Pharmacology. $2.75

COBLENTZ. The Newer Remedies. Including their Synonyms, Sources, Methods of Preparation, Tests, Solubilities, Doses, etc. 3d Edition, Enlarged and Revised. $1.00

COHEN. Physiologic Therapeutics. Mechanotherapy, Mental Therapeutics, Electrotherapy, Climatology, Hydrotherapy, Pneumotherapy, Prophylaxis, Alimentotherapy, etc. *In Press.*

DAVIS. Materia Medica and Prescription Writing. $1.50

GORGAS. Dental Medicine. A Manual of Materia Medica and Therapeutics. 6th Edition, Revised. $4.00

GROFF. Materia Medica for Nurses, with questions for Self Examination and a complete Glossary. $1.25

HELLER. Essentials of Materia Medica, Pharmacy, and Prescription Writing. $1.50

MAYS. Theine in the Treatment of Neuralgia. ½ bound, .50

POTTER. Hand-Book of Materia Medica, Pharmacy, and Therapeutics, including the Action of Medicines, Special Therapeutics, Pharmacology, etc., including over 600 Prescriptions and Formulæ. 7th Edition, Revised and Enlarged. With Thumb Index in each copy. Cloth, $5.00; Sheep, $6.00

POTTER. Compend of Materia Medica, Therapeutics, and Prescription Writing, with Special Reference to the Physiological Action of Drugs. 6th Edition. .80; Interleaved, $1.00

SAYRE. Organic Materia Medica and Pharmacognosy. An Introduction to the Study of the Vegetable Kingdom and the Vegetable and Animal Drugs. Comprising the Botanical and Physical Characteristics, Source, Constituents, and Pharmacopeial Preparations, Insects Injurious to Drugs, and Pharmacal Botany. With sections on Histology and Microtechnique, by W. C. STEVENS. 374 Illustrations, many of which are original. 2d Edition.
Cloth, $4.50

WHITE AND WILCOX. Materia Medica, Pharmacy, Pharmacology, and Therapeutics. 4th American Edition, Revised by REYNOLD W. WILCOX, M.A., M D., LL.D., Professor of Clinical Medicine and Therapeutics at the New York Post-Graduate Medical School. Cloth, $3 00; Leather, $3.50

"The care with which Dr. Wilcox has performed his work is conspicuous on every page, and it is evident that no recent drug possessing any merit has escaped his eye. We believe, on the whole, this is the best book on Materia Medica and Therapeutics to place in the hands of students, and the practitioner will find it a most satisfactory work for daily use."—*The Cleveland Medical Gazette.*

MEDICAL JURISPRUDENCE AND TOXICOLOGY.

REESE. Medical Jurisprudence and Toxicology. A Text-Book for Medical and Legal Practitioners and Students. 5th Edition. Revised by HENRY LEFFMANN, M.D. Clo., $3.00; Leather, $3.50

"To the student of medical jurisprudence and toxicology it is invaluable, as it is concise, clear, and thorough in every respect."—*The American Journal of the Medical Sciences.*

MANN. Forensic Medicine and Toxicology. Illus. $6.50

TANNER. Memoranda of Poisons. Their Antidotes and Tests. 7th Edition. .75

MICROSCOPY.

CARPENTER. The Microscope and Its Revelations. 8th Edition. 800 Illustrations and many Lithographs. *Preparing.*

LEE. The Microtomist's Vade Mecum. A Hand-Book of Methods of Microscopical Anatomy. 887 Articles. 5th Edition, Enlarged. *Just Ready.* $4.00

REEVES. Medical Microscopy, including Chapters on Bacteriology, Neoplasms, Urinary Examination, etc. Numerous Illustrations, some of which are printed in colors. $2.50

WETHERED. Medical Microscopy. A Guide to the Use of the Microscope in Practical Medicine. 100 Illustrations. $2.00

MISCELLANEOUS.

BRAMWELL. Anæmia. $2.50
BURNETT. Foods and Dietaries. A Manual of Clinical Dietetics. 2d Edition. $1.50
BUXTON. Anæsthetics. Illustrated. 3d Edition. $1.50
COHEN. Organotherapy. *In Press.*
DAVIS. Alimentotherapy. *In Press.*
FENWICK. Ulcer of the Stomach. 42 Illustrations. $3.50
GOULD. Borderland Studies. Miscellaneous Addresses and Essays. 12mo. $2.00
GREENE. Medical Examination for Life Insurance. Illustrated. $1.00
HAIG. Causation of Disease by Uric Acid. The Pathology of High Arterial Tension, Headache, Epilepsy, Gout, Rheumatism, Diabetes, Bright's Disease, etc. 5th Edition. *Just Ready.* $3.00
HAIG. Diet and Food. Considered in Relation to Strength and Power of Endurance. 2d Edition. $1.00
HEMMETER. Diseases of the Stomach. Their Special Pathology, Diagnosis, and Treatment. With Sections on Anatomy, Dietetics, Surgery, etc. 2d Edition, Revised and Enlarged. Illustrated. *Just Ready.* Cloth, $6.00; Sheep, $7.00
HEMMETER. Intestines. Illustrated. *Preparing.*
HENRY. A Practical Treatise on Anemia. Half Cloth, .50
MARSHALL. History of Woman's Medical College of Pennsylvania. $1.50
MATAS. Anesthesia with Cocain, Eucain, etc. Local and Regional. Illustrated. *Nearly Ready.*
NEW SYDENHAM SOCIETY'S PUBLICATIONS. Circulars upon application. Per Annum, $8.00
OSGOOD. The Winter and Its Dangers. .40
OSLER AND McCRAE. Cancer of the Stomach. *Just Ready.* $2.00
PACKARD. Sea Air and Sea Bathing. .40
RICHARDSON. Long Life and How to Reach It. .40
ST. CLAIR. Medical Latin. $1.00
TESSIER. Pneumotherapy. *In Press.*
TREVES. Physical Education: Its Effects, Methods, Etc. .75
TURNBULL. Artificial Anæsthesia. 4th Edition. Illus. $2.50
WEBER AND HINSDALE. Climatology. *In Press.*
WILSON. The Summer and Its Diseases. .40
WINTERNITZ. Hydrotherapy. *In Press.*

NERVOUS DISEASES.

BEEVOR. Diseases of the Nervous System and their Treatment. $2.50
DERCUM. Rest, Hypmotism, Mental Therapeutics. *In Press.*
GORDINIER. The Gross and Minute Anatomy of the Central Nervous System. With 271 original Colored and other Illustrations. Cloth, $6.00; Sheep, $7.00
GOWERS. Manual of Diseases of the Nervous System. A Complete Text-Book. Revised, Enlarged, and in many parts Rewritten. With many new Illustrations. Two volumes.
Vol. I. Diseases of the Nerves and Spinal Cord. 3d Edition, Enlarged. Cloth, $4.00; Sheep, $5.00
Vol. II. Diseases of the Brain and Cranial Nerves: General and Functional Diseases. 2d Edition. Cloth, $4.00; Sheep, $5.00
GOWERS. Syphilis and the Nervous System. $1.00

GOWERS. Clinical Lectures. A New Volume of Essays on the Diagnosis, Treatment, etc., of Diseases of the Nervous System. $2.00

GOWERS. Epilepsy and Other Chronic Convulsive Diseases. 2d Edition. *In Press.*

HORSLEY. The Brain and Spinal Cord. The Structure and Functions of. Numerous Illustrations. $2.50

ORMEROD. Diseases of the Nervous System. 66 Wood Engravings. $1.00

OSLER. Chorea and Choreiform Affections. $2.00

PRESTON. Hysteria and Certain Allied Conditions. Their Nature and Treatment. Illustrated. $2.00

WOOD. Brain Work and Overwork. .40

NURSING (see also Massage).

Special Catalogue of Books for Nurses sent free upon application.

BROWN. Elementary Physiology for Nurses. .75

CANFIELD. Hygiene of the Sick-Room. A Book for Nurses and Others. Being a Brief Consideration of Asepsis, Antisepsis, Disinfection, Bacteriology, Immunity, Heating and Ventilation, and Kindred Subjects for the Use of Nurses and Other Intelligent Women. $1.25

CUFF. Lectures to Nurses on Medicine. New Edition. $1.25

DOMVILLE. Manual for Nurses and Others Engaged in Attending the Sick. 8th Edition. With Recipes for Sick-room Cookery, etc. .75

FULLERTON. Obstetric Nursing. 41 Ills. 5th Ed. $1.00

FULLERTON. Surgical Nursing. 3d Ed. 69 Ills. $1.00

GROFF. Materia Medica for Nurses. With Questions for Self-Examination and a very complete Glossary. $1.25

"It will undoubtedly prove a valuable aid to the nurse in securing a knowledge of drugs and their uses."—*The Medical Record, New York.*

HORWITZ. Duties of the Surgical Nurse. *In Press.*

HUMPHREY. A Manual for Nurses. Including General Anatomy and Physiology, Management of the Sick Room, etc. 17th Ed. Illustrated. $1.00

"In the fullest sense, Dr. Humphrey's book is a distinct advance on all previous manuals. It is, in point of fact, a concise treatise on medicine and surgery for the beginner, incorporating with the text the management of childbed and the hygiene of the sick-room. Its value is greatly enhanced by copious wood-cuts and diagrams of the bones and internal organs."—*British Medical Journal, London.*

STARR. The Hygiene of the Nursery. Including the General Regimen and Feeding of Infants and Children, and the Domestic Management of the Ordinary Emergencies of Early Life, Massage, etc. 6th Edition. 25 Illustrations. $1.00

TEMPERATURE AND CLINICAL CHARTS. See page 6.

VOSWINKEL. Surgical Nursing. Second Edition, Enlarged. 112 Illustrations. $1.00

WESTLAND. The Wife and Mother. $1.50

OBSTETRICS.

CAZEAUX AND TARNIER. Midwifery. With Appendix by MUNDÉ. The Theory and Practice of Obstetrics, including the Diseases of Pregnancy and Parturition, Obstetrical Operations, etc. 8th Edition. Illustrated by Chromo-Lithographs, Lithographs, and other full-page Plates, seven of which are beautifully colored, and numerous Wood Engravings. Cloth, $4.50; Full Leather, $5.50

DAVIS. A Manual of Obstetrics. 3d Edition. *Preparing.*

EDGAR. Text-Book of Obstetrics. Illustrated. *Preparing.*

LANDIS. Compend of Obstetrics. 6th Edition, Revised by WM. H. WELLS, Assistant Demonstrator of Clinical Obstetrics, Jefferson Medical College. With 47 Illustrations, .80; Interleaved, $1.00

WINCKEL. Text-Book of Obstetrics, Including the Pathology and Therapeutics of the Puerperal State. Authorized Translation by J. CLIFTON EDGAR, A.M., M.D. With nearly 200 Illustrations. Cloth, $5.00; Leather, $6.00

FULLERTON. Obstetric Nursing. 5th Ed. Illustrated. $1.00

PATHOLOGY.

BARLOW. General Pathology. 795 pages. 8vo. $5.00

BLACK. Micro-Organisms. The Formation of Poisons. .75

BLACKBURN. Autopsies. A Manual of Autopsies Designed for the Use of Hospitals for the Insane and other Public Institutions. Ten full-page Plates and other Illustrations. $1.25

COMPEND of General Pathology. Illustrated. *Preparing.*

COPLIN. Manual of Pathology. Including Bacteriology, Technic of Post-Mortems, Methods of Pathologic Research, etc. 330 Illustrations, 7 Colored Plates, many of which are original. 3d Edition. *Just Ready.* $3.50

DA COSTA. Clinical Pathology of the Blood. Illus. *In Press.*

GILLIAM. Pathology. A Hand-Book for Students. 47 Illus. .75

HEWLETT. Manual of Bacteriology. 75 Illustrations. $3.00

VIRCHOW. Post-Mortem Examinations. A Description and Explanation of the Method of Performing Them in the Dead House of the Berlin Charity Hospital, with Special Reference to Medico-Legal Practice. 3d Edition, with Additions. .75

WHITACRE. Laboratory Text-Book of Pathology. With 121 Illustrations. $1.50

WILLIAMS. Bacteriology. A Manual for Students. 78 Illustrations. $1.50

PHARMACY.

Special Catalogue of Books on Pharmacy sent free upon application.

COBLENTZ. Manual of Pharmacy. A Complete Text-Book by the Professor in the New York College of Pharmacy. 2d Edition, Revised and Enlarged. 437 Illus. Cloth, $3.50; Sheep, $4.50

BEASLEY. Book of 3100 Prescriptions. Collected from the Practice of the Most Eminent Physicians and Surgeons—English, French, and American. A Compendious History of the Materia Medica, Lists of the Doses of all the Officinal and Established Preparations, an Index of Diseases and their Remedies. 7th Ed. $2.00

BEASLEY. Druggists' General Receipt Book. Comprising a Copious Veterinary Formulary, Recipes in Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics, Beverages, Dietetic Articles and Condiments, Trade Chemicals, Scientific Processes, and many Useful Tables. 10th Ed. $2.00

BEASLEY. Pharmaceutical Formulary. A Synopsis of the British, French, German, and United States Pharmacopœias. Comprising Standard and Approved Formulæ for the Preparations and Compounds Employed in Medicine. 12th Edition. $2.00

PROCTOR. Practical Pharmacy. Lectures on Practical Pharmacy. With Wood Engravings and 32 Lithographic Fac-simile Prescriptions. 3d Edition, Revised, and with Elaborate Tables of Chemical Solubilities, etc. $3.00

ROBINSON. Latin Grammar of Pharmacy and Medicine. 3d Edition. With elaborate Vocabularies. $1.75

SAYRE. Organic Materia Medica and Pharmacognosy. An Introduction to the Study of the Vegetable Kingdom and the Vegetable and Animal Drugs. Comprising the Botanical and Physical Characteristics, Source, Constituents, and Pharmacopeial Preparations, Insects Injurious to Drugs, and Parmacal Botany. With sections on Histology and Microtechnique, by W. C. Stevens. 374 Illustrations. Second Edition. Cloth, $4.50

SCOVILLE. The Art of Compounding. Second Edition, Revised and Enlarged. Cloth, $2.50

STEWART. Compend of Pharmacy. Based upon "Remington's Text-Book of Pharmacy." 5th Edition, Revised in Accordance with the U. S. Pharmacopœia, 1890. Complete Tables of Metric and English Weights and Measures. .80; Interleaved, $1.00

UNITED STATES PHARMACOPŒIA. 7th Decennial Revision. Cloth, $2.50 (postpaid, $2.77); Sheep, $3.00 (postpaid, $3.27); Interleaved, $4.00 (postpaid, $4.50); Printed on one side of page only, unbound, $3.50 (postpaid, $3.90).

Select Tables from the U. S. P. Being Nine of the Most Important and Useful Tables, Printed on Separate Sheets. Carefully put up in patent envelope. .25

POTTER. Hand-Book of Materia Medica, Pharmacy, and Therapeutics. 600 Prescriptions. 7th Ed. Clo., $5.00; Sh., $6.00

PHYSICAL DIAGNOSIS.

BROWN. Medical Diagnosis. A Manual of Clinical Methods. 4th Edition. 112 Illustrations. Cloth, $2.25

DA COSTA. Clinical Examination of the Blood. Illustrated. *In Press.*

FENWICK. Medical Diagnosis. 8th Edition. Rewritten and very much Enlarged. 135 Illustrations. Cloth, $2.50

MEMMINGER. Diagnosis by the Urine. 2d Ed. 24 Illus. $1 00

TYSON. Hand-Book of Physical Diagnosis. For Students and Physicians. By the Professor of Clinical Medicine in the University of Pennsylvania. Illus. 3d Ed., Improved and Enlarged. With Colored and other Illustrations. $1.50

PHYSIOLOGY.

BIRCH. Practical Physiology. An Elementary Class Book. 62 Illustrations. $1.75

BRUBAKER. Compend of Physiology. 10th Edition, Revised and Enlarged. Illustrated. *Just Ready.* .80; Interleaved, $1.00

KIRKES. Physiology. (16th Authorized Edition. Dark-Red Cloth.) A Hand-Book of Physiology. 16th Edition, Revised, Rearranged, and Enlarged. By PROF. W. D. HALLIBURTON, of Kings College, London. 671 Illustrations, some of which are printed in colors. *Just Ready.* Cloth, $3.00; Leather, $3.75

LANDOIS. A Text-Book of Human Physiology, Including Histology and Microscopical Anatomy, with Special Reference to the Requirements of Practical Medicine. 5th American, translated from the 9th German Edition, with Additions by WM. STIRLING, M.D.,D.SC. 845 Illus., many of which are printed in colors. *In Press.*

STARLING. Elements of Human Physiology. 100 Ills. $1.00

STIRLING. Outlines of Practical Physiology. Including Chemical and Experimental Physiology, with Special Reference to Practical Medicine. 3d Edition. 289 Illustrations. $2.00

TYSON. Cell Doctrine. Its History and Present State. $1.50

PRACTICE.

BEALE. On Slight Ailments; their Nature and Treatment. 2d Edition, Enlarged and Illustrated. $1.25

FOWLER. Dictionary of Practical Medicine. By various writers. An Encyclopædia of Medicine. Clo., $3.00; Half Mor. $4.00

GOULD AND PYLE. Cyclopedia of Practical Medicine and Surgery. A Concise Reference Handbook, Alphabetically Arranged, with particular Reference to Diagnosis and Treatment. Edited by DRS. GOULD and PYLE, Assisted by 72 Special Contributors. Illustrated, one volume. Large Square Octavo, Uniform with "Gould's Illustrated Dictionary." *Just Ready.*

Sheep or Half Morocco, $10.00; with Thumb Index, $11.00
Half Russia, Thumb Index, $12 00

☞ *Complete descriptive circular with sample pages and illustrations of this book will be sent free upon application.*

HUGHES. Compend of the Practice of Medicine. 6th Edition, Revised and Enlarged. *Just Ready.*

Part I. Continued, Eruptive, and Periodical Fevers, Diseases of the Stomach, Intestines, Peritoneum, Biliary Passages, Liver, Kidneys, etc., and General Diseases, etc.

Part II. Diseases of the Respiratory System, Circulatory System, and Nervous System; Diseases of the Blood, etc.

Price of each part, .80; Interleaved, $1.00

Physician's Edition. In one volume, including the above two parts, a Section on Skin Diseases, and an Index. 6th Revised Edition. 625 pp. *Just Ready.* Full Morocco, Gilt Edge, $2.25

TAYLOR. Practice of Medicine. 5th Edition. Cloth, $4.00

TYSON. The Practice of Medicine. By JAMES TYSON, M.D., Professor of Medicine in the University of Pennsylvania. A Complete Systematic Text-book with Special Reference to Diagnosis and Treatment. 2d Edition, Enlarged and Revised. Colored Plates and 125 other Illustrations. 1222 Pages. 8vo. *Just Ready.*

Cloth, $5.50; Leather, $6.50

PRESCRIPTION BOOKS.

BEASLEY. Book of 3100 Prescriptions. Collected from the Practice of the Most Eminent Physicians and Surgeons—English, French, and American. A Compendious History of the Materia, Medica, Lists of the Doses of all Officinal and Established Preparations, and an Index of Diseases and their Remedies. 7th Ed. $2.00

BEASLEY. Druggists' General Receipt Book. Comprising a Copious Veterinary Formulary, Recipes in Patent and Proprietary Medicines, Druggists' Nostrums, etc.; Perfumery and Cosmetics, Beverages, Dietetic Articles and Condiments, Trade Chemicals, Scientific Processes, and an Appendix of Useful Tables. 10th Edition, Revised. $2.00

BEASLEY. Pocket Formulary. A Synopsis of the British, French, German, and United States Pharmacopœias and the chief unofficial Formularies. 12th Edition. $2.00

SKIN.

BULKLEY. The Skin in Health and Disease. Illustrated. .40

CROCKER. Diseases of the Skin. Their Description, Pathology, Diagnosis, and Treatment, with Special Reference to the Skin Eruptions of Children. 92 Illus. 3d Edition. *Preparing.*

IMPEY. Leprosy. 37 Plates. 8vo. $3.50

SCHAMBERG. Diseases of the Skin. 2d Edition, Revised and Enlarged. 105 Illustrations. Being No. 16 ? Quiz-Compend ? Series. *Just Ready.* Cloth, .80; Interleaved, $1.00

VAN HARLINGEN. On Skin Diseases. A Practical Manual of Diagnosis and Treatment, with special reference to Differential Diagnosis. 3d Edition, Revised and Enlarged. With Formulæ and 60 Illustrations, some of which are printed in colors. $2.75

SURGERY AND SURGICAL DISEASES (see also Urinary Organs).

BUTLIN. Operative Surgery of Malignant Disease. 2d Edition. Illustrated. Octavo. *Just Ready.* $4.50

CRIPPS. Ovariotomy and Abdominal Surgery. Illus. $8.00

DEAVER. Surgical Anatomy. A Treatise on Human Anatomy in its Application to Medicine and Surgery. With about 400 very Handsome full-page Illustrations Engraved from Original Drawings made by special Artists from Dissections prepared for the purpose. Three Volumes. Royal Square Octavo.
Cloth, $21.00; Half Morocco or Sheep, $24.00; Half Russia, $27.00
Complete descriptive circular and special terms upon application.

DEAVER. Appendicitis, Its Symptoms, Diagnosis, Pathology, Treatment, and Complications. Elaborately Illustrated with Colored Plates and other Illustrations. 2d Edition. $3.50

DULLES. What to Do First in Accidents and Poisoning. 5th Edition. New Illustrations. $1.00

FULLERTON. Surgical Nursing. 3d Edition. 69 Illus. $1 00

HAMILTON. Lectures on Tumors, from a Clinical Standpoint. Third Edition, Revised, with New Illustrations. $1.25

HEATH. Minor Surgery and Bandaging. 11th Ed., Revised and Enlarged. 176 Illustrations, Formulæ, Diet List, etc. $1.25

HEATH. Injuries and Diseases of the Jaws. 4th Edition. 187 Illustrations. $4.50

HEATH. Lectures on Certain Diseases of the Jaws. 64 Illustrations. Boards, .50

HORWITZ. Compend of Surgery and Bandaging, including Minor Surgery, Amputations, Fractures, Dislocations, Surgical Diseases, and the Latest Antiseptic Rules, etc., with Differential Diagnosis and Treatment. 5th Edition, very much Enlarged and Rearranged. 167 Illustrations, 98 Formulæ. Clo., .80; Interleaved, $1.25

JACOBSON. Operations of Surgery. Over 200 Illustrations. Cloth, $3.00; Leather, $4.00

JACOBSON. Diseases of the Male Organs of Generation. 88 Illustrations. $6.00

LANE. Surgery of the Head and Neck. 110 Illustrations. 2d Edition. $5.00

MACREADY. A Treatise on Ruptures. 24 Full-page Lithographed Plates and Numerous Wood Engravings. Cloth, $6.00

MAYLARD. Surgery of the Alimentary Canal. 97 Illustrations. 2d Edition, Revised. *Just Ready.* $3.00

MOULLIN. Text-Book of Surgery. With Special Reference to Treatment. 3d American Edition. Revised and edited by JOHN B. HAMILTON, M.D., LL.D., Professor of the Principles of Surgery and Clinical Surgery, Rush Medical College, Chicago. 623 Illustrations, over 200 of which are original, and many of which are printed in colors. Handsome Cloth, $6.00; Leather, $7.00

ROBERTS. Fractures of the Radius. A Clinical and Pathological Study. 33 Illustrations. $1.00

SMITH. Abdominal Surgery. Being a Systematic Description of all the Principal Operations. 224 Illus. 6th Ed. 2 Vols. Clo., $10.00

SWAIN. Surgical Emergencies. Fifth Edition. Cloth, $1.75

VOSWINKEL. Surgical Nursing. Second Edition, Revised and Enlarged. 111 Illustrations. $1.00

WALSHAM. Manual of Practical Surgery. 7th Ed., Revised and Enlarged. 483 Engravings. 950 pages. *Just Ready.* $3.50

THROAT AND NOSE (see also Ear).

COHEN. The Throat and Voice. Illustrated. .40

HALL. Diseases of the Nose and Throat. Two Colored Plates and 59 Illustrations. New Edition Preparing.

HOLLOPETER. Hay Fever. Its Successful Treatment. $1.00

KNIGHT. Diseases of the Throat. A Manual for Students. Illustrated. *Nearly Ready.*

MACKENZIE. Pharmacopœia of the London Hospital for Dis. of the Throat. 5th Ed., Revised by Dr. F. G. HARVEY. $1.00

McBRIDE. Diseases of the Throat, Nose, and Ear. With colored Illus. from original drawings. 3d Ed. *Just Ready* $7.00

POTTER. Speech and its Defects. Considered Physiologically, Pathologically, and Remedially. $1.00

URINE AND URINARY ORGANS.

ACTON. The Functions and Disorders of the Reproductive Organs in Childhood, Youth, Adult Age, and Advanced Life, Considered in their Physiological, Social, and Moral Relations. 8th Edition. $1.75

BEALE. One Hundred Urinary Deposits. On eight sheets, for the Hospital, Laboratory, or Surgery. Paper, $2.00

HOLLAND. The Urine, the Gastric Contents, the Common Poisons, and the Milk. Memoranda, Chemical and Microscopical, for Laboratory Use. Illustrated and Interleaved. 6th Ed. $1.00

JACOBSON. Male Organs of Generation. 88 Illus. $6.00

KEHR. Gall-Stone Disease. Translated by WILLIAM WATKYNS SEYMOUR, M D. *In Press.*

KLEEN. Diabetes and Glycosuria. $2.50

MEMMINGER. Diagnosis by the Urine. 2d Ed. 24 Illus. $1.00

MORRIS. Renal Surgery, with Special Reference to Stone in the Kidney and Ureter and to the Surgical Treatment of Calculous Anuria. Illustrated. $2.00.

MOULLIN. Enlargement of the Prostate. Its Treatment and Radical Cure. 2d Edition. Illustrated. *Just Ready.* $1.75

MOULLIN. Inflammation of the Bladder and Urinary Fever. Octavo. $1.50

SCOTT. The Urine. Its Clinical and Microscopical Examination. 41 Lithographic Plates and other Illustrations. Quarto. Cloth, $5.00

TYSON. Guide to Examination of the Urine. For the Use of Physicians and Students. With Colored Plate and Numerous Illustrations engraved on wood. 9th Edition, Revised. $1.25

VAN NUYS. Chemical Analysis of Urine. 39 Illus. $1.00

VENEREAL DISEASES.

COOPER. Syphilis. 2d Edition, Enlarged and Illustrated with 20 full-page Plates. $5.00

GOWERS. Syphilis and the Nervous System. 1.00

STURGIS. Student's Manual of Venereal Diseases. 7th Revised and Enlarged Edition. 12mo. *In Press.*

VETERINARY.

BALLOU. Veterinary Anatomy and Physiology. 29 Graphic Illustrations. .80; Interleaved, $1.00

TUSON. Veterinary Pharmacopœia. Including the Outlines of Materia Medica and Therapeutics. 5th Edition. $2.25

WOMEN, DISEASES OF.

BYFORD (H. T.). Manual of Gynecology. Second Edition, Revised and Enlarged by 100 pages. 341 Illustrations. $3.00

DÜHRSSEN. A Manual of Gynecological Practice. 105 Illustrations. $1.50

FULLERTON. Surgical Nursing. 3d Edition, Revised and Enlarged. 69 Illustrations. $1.00

LEWERS. Diseases of Women. 146 Illus. 5th Ed. $2.50

MONTGOMERY. Practical Gynecology. A Complete Systematic Text-Book. 527 Handsome Illustrations. 8vo. *Just Ready.* Cloth, $5.00; Leather, $6.00

WELLS. Compend of Gynecology. Illustrated. 2d Edition. .80; Interleaved, $1.00

COMPENDS.

From The Southern Clinic.

"We know of no series of books issued by any house that so fully meets our approval as these ? Quiz-Compends ?. They are well arranged, full, and concise, and are really the best line of text-books that could be found for either student or practitioner."

BLAKISTON'S ? QUIZ-COMPENDS?

The Best Series of Manuals for the Use of Students.

Price of each, Cloth, .80. Interleaved, for taking Notes, $1.00.

☞ These Compends are based on the most popular text-books and the lectures of prominent professors, and are kept constantly revised, so that they may thoroughly represent the present state of the subjects upon which they treat.

☞ The authors have had large experience as Quiz-Masters and attaches of colleges, and are well acquainted with the wants of students.

☞ They are arranged in the most approved form, thorough and concise, containing over 600 fine illustrations, inserted wherever they could be used to advantage.

☞ Can be used by students of *any* college.

☞ They contain information nowhere else collected in such a condensed, practical shape. **Illustrated Circular free.**

No. 1. POTTER. HUMAN ANATOMY. Sixth Revised and Enlarged Edition. Including Visceral Anatomy. Can be used with either Morris's or Gray's Anatomy. 117 Illustrations and 16 Lithographic Plates of Nerves and Arteries, with Explanatory Tables, etc. By SAMUEL O. L. POTTER, M.D., Professor of the Practice of Medicine, College of Physicians and Surgeons, San Francisco; Brigade Surgeon, U. S. Vol.

No. 2. HUGHES. PRACTICE OF MEDICINE. Part I. Sixth Edition, Enlarged and Improved. By DANIEL E. HUGHES, M.D., Physician-in-Chief, Philadelphia Hospital, late Demonstrator of Clinical Medicine, Jefferson Medical College, Phila.

No. 3. HUGHES. PRACTICE OF MEDICINE. Part II. Sixth Edition, Revised and Improved. Same author as No. 2.

No. 4. BRUBAKER. PHYSIOLOGY. Tenth Edition, with Illustrations and a table of Physiological Constants. Enlarged and Revised. By A. P. BRUBAKER, M.D., Professor of Physiology and General Pathology in the Pennsylvania College of Dental Surgery; Adjunct Professor of Physiology, Jefferson Medical College, Philadelphia, etc

No. 5. LANDIS. OBSTETRICS. Sixth Edition. By HENRY G. LANDIS, M.D. Revised and Edited by WM. H. WELLS, M.D., Instructor of Obstetrics, Jefferson Medical College, Philadelphia. Enlarged. 47 Illustrations.

No. 6. POTTER. MATERIA MEDICA, THERAPEUTICS, AND PRESCRIPTION WRITING. Sixth Revised Edition (U. S. P. 1890). By SAMUEL O. L. POTTER, M.D., Professor of Practice, College of Physicians and Surgeons, San Francisco; Brigade Surgeon, U. S. Vol.

? QUIZ-COMPENDS ?—Continued.

No. 7. WELLS. GYNECOLOGY. Second Edition. By WM. H. WELLS, M.D., Instructor of Obstetrics, Jefferson College, Philadelphia. 140 Illustrations.

No. 8. GOULD AND PYLE. DISEASES OF THE EYE AND REFRACTION. Second Edition. Including Treatment and Surgery, and a Section on Local Therapeutics. By GEORGE M. GOULD, M.D., and W. L. PYLE, M.D. With Formulæ, Glossary, Tables, and 109 Illustrations, several of which are Colored.

No. 9. HORWITZ. SURGERY, Minor Surgery, and Bandaging. Fifth Edition, Enlarged and Improved. By ORVILLE HORWITZ, B.S., M.D., Clinical Professor of Genito-Urinary Surgery and Venereal Diseases in Jefferson Medical College; Surgeon to Philadelphia Hospital, etc. With 98 Formulæ and 71 Illustrations.

No. 10. LEFFMANN. MEDICAL CHEMISTRY. Fourth Edition. Including Urinalysis, Animal Chemistry, Chemistry of Milk, Blood, Tissues, the Secretions, etc. By HENRY LEFFMANN, M.D., Professor of Chemistry in Pennsylvania College of Dental Surgery and in the Woman's Medical College, Philadelphia.

No. 11. STEWART. PHARMACY. Fifth Edition. Based upon Prof. Remington's Text-Book of Pharmacy. By F. E. STEWART, M.D., PH.G., late Quiz-Master in Pharmacy and Chemistry, Philadelphia College of Pharmacy; Lecturer at Jefferson Medical College. Carefully revised in accordance with the new U. S. P.

No. 12. BALLOU. VETERINARY ANATOMY AND PHYSIOLOGY. Illustrated. By WM. R. BALLOU, M.D., Professor of Equine Anatomy at New York College of Veterinary Surgeons; Physician to Bellevue Dispensary, etc. 29 graphic Illustrations

No. 13. WARREN. DENTAL PATHOLOGY AND DENTAL MEDICINE. Third Edition, Illustrated. Containing a Section on Emergencies. By GEO. W. WARREN, D.D.S., Chief of Clinical Staff, Pennsylvania College of Dental Surgery.

No. 14. HATFIELD. DISEASES OF CHILDREN. Second Edition. **Colored Plate.** By MARCUS P. HATFIELD, Professor of Diseases of Children, Chicago Medical College.

No. 15. GENERAL PATHOLOGY. Illustrated *Preparing.*

No. 16. DISEASES OF THE SKIN. Second Edition. By JAY F. SCHAMBERG, M.D., Professor of Diseases of the Skin, Philadelphia Polyclinic. Second Edition, Revised and Enlarged. 105 handsome Illustrations.

Price, each, Cloth, .80. Interleaved, for taking Notes, $1.00

In preparing, revising, and improving BLAKISTON'S ? QUIZ-COMPENDS ? the particular wants of the student have always been kept in mind.

Careful attention has been given to the construction of each sentence, and while the books will be found to contain an immense amount of knowledge in small space, they will likewise be found easy reading; there is no stilted repetition of words; the style is clear, lucid, and distinct. The arrangement of subjects is systematic and thorough; there is a reason for every word. They contain over 600 illustrations.

Morris' Anatomy

Second Edition, Revised and Enlarged.

790 Illustrations, of which many are in Colors.

Royal Octavo. Cloth, $6.00; Sheep, $7.00; Half Russia, $8.00.

From The Medical Record, New York.

"The reproach that the English language can boast of no treatise on anatomy deserving to be ranked with the masterly works of Henle, Luschka, Hyrtl, and others, is fast losing its force. During the past few years several works of great merit have appeared, and among these Morris's "Anatomy" seems destined to take first place in disputing the palm in anatomical fields with the German classics. The nomenclature, arrangement, and entire general character resemble strongly those of the above-mentioned handbooks, while in the beauty and profuseness of its illustrations it surpasses them. . . . The ever-growing popularity of the book with teachers and students is an index of its value, and it may safely be recommended to all interested."

From The Philadelphia Medical Journal.

"Of all the text-books of moderate size on human anatomy in the English language, Morris is undoubtedly the most up-to-date and accurate."

***** Handsome Descriptive Circular, with Sample Pages and Colored Illustrations, will be sent free upon application.**

28.E.39.
Retinoscopy (or shadow test) in1901
Countway Library BEY8641
3 2044 046 138 897

28.E.39.
Retinoscopy (or shadow test) in1901
Countway Library BEY8641
3 2044 046 138 897